

Title: Quantum Field Theory for Cosmology - Lecture 11

Speakers: Achim Kempf

Collection: Quantum Field Theory for Cosmology (Kempf)

Date: February 11, 2020 - 4:00 PM

URL: <http://pirsa.org/20020002>

QFT for Cosmology, Achim Kempf, Lecture 11

Recall the strategy:

SR, 1st Q
Hamiltonian
formalism

step 1
Legendre transform
(equivalence)

SR, 1st Q.
Lagrangian
formalism

step 2
allow
curvature

GR, 1st Q
Hamiltonian
formalism

step 3
Legendre transform
(equivalence)

GR, 1st Q
Lagrangian
formalism

step 4
GR, 2nd Q
Hamiltonian
formalism

Dyson Schwinger eqns are same
(equivalence)

GR, 2nd Q
Lagrangian formalism
(Path integral of QFT)



Type here to search



4:02 PM
ENG
2020-02-11

Curvature:

- We postulate the coordinate system-independent Klein-Gordon action:

$$S[\phi] := \frac{1}{2} \int_{\text{M}} \left(g^{\mu\nu} \partial_\mu \phi_{,\nu} - m^2 \phi^2 - V(\phi, x_i) \right) \sqrt{|g|} d^4x$$

- We will allow almost arbitrary metric tensors $g_{\mu\nu}(x)$, even those for which there do not exist coordinates \tilde{x} in which:

$$\tilde{g}_{\mu\nu}(\tilde{x}) = \eta_{\mu\nu} \quad \text{for all } x$$

- But we must have that, at least locally, special relativity holds!

⇒ Consider only $g_{\mu\nu}(x)$ for which for each x_0 , there exists a change of coordinates $x \rightarrow \tilde{x}$

so that:

$$\tilde{g}_{\mu\nu}(\tilde{x}_0) = \eta_{\mu\nu}$$

* We assume that if a freely falling observer in a small region sets up a rectangular coordinate system the observer will see arbitrarily small gravity effects if the region is made arbitrarily small.

* For this to be true, any body's gravitational mass must be equal to its inertial mass, i.e. all bodies must fall equally. (Else the notion of freely falling observer is not even well defined)

How can we identify the presence of curvature?



Type here to search

4:06 PM
ENG
2020-02-11

How can one identify the presence of curvature?

* Assume we are given a metric tensor

$$g_{\mu\nu}(x)$$


as an explicit matrix-valued function, in some coordinates.

* How can we determine whether or not this is, e.g., the metric tensor of flat space-time, i.e., whether or not there exist coordinates \tilde{x} in which $\tilde{g}_{\mu\nu}(\tilde{x}) = g_{\mu\nu}$?

* This problem is solved in differential geometry:

Define: The "Christoffel symbol functions":

$$\Gamma_{\alpha\beta}^{\rho}(x) := \frac{1}{2} g^{\rho\sigma}(x) \left(g_{\alpha\sigma,\beta}(x) + g_{\beta\sigma,\alpha}(x) - g_{\alpha\beta,\sigma}(x) \right)$$




Define: The "Riemann Curvature Tensor":

$$R^i_{jkl}(x) := \Gamma^i_{ljk}(x) - \Gamma^i_{kjl}(x) + \Gamma^s_{lj}(x)\Gamma^i_{ks}(x) - \Gamma^s_{kj}(x)\Gamma^i_{ls}(x)$$



It's rôle? A space is called curved at x if the parallel transport of a vector v along an infinitesimal parallelogram returns the vector v' to x , but v' is rotated by some amount. R^r_{vpl} tells by how much:



$$(v' - v)^a = \eta^b \eta^c R^a_{bcd}(x) v^d$$



Type here to search



4:08 PM ENG 2020-02-11

Remark:

If the parallelogram does not even close we say that space-time has "Torsion". There is no evidence for torsion in nature.

Proposition:

Assume that, in a region, A , of space-time :

$$R^{\mu}_{\nu\lambda\rho}(x) = 0 \text{ for all } x \in A$$

Then and only then there exist coordinates \tilde{x} so that:

$$\tilde{g}_{\mu\nu}(\tilde{x}) = g_{\mu\nu} \text{ for all } x \in A$$



The dynamics of space-time

Problem: What are the equations of motion for spacetime's curvature?

Which are the degrees of freedom of curvature
for which we have to find an equation of motion?

□ We saw that the curvature of space-time is
encoded in the matrix-valued metric function:

$$g_{\mu\nu}(x)$$

□ However, if $g_{\mu\nu}(x)$ looks nontrivial, this can
be for two different reasons:

1. Spacetime has little or no curvature and $g_{\mu\nu}(x)$ is nontrivial just because of an unlucky choice of coordinates.

2. Spacetime is curved, i.e., we can not make $g_{\mu\nu}(x)$ take the form $\tilde{g}_{\mu\nu}(\hat{x}) = \eta_{\mu\nu}$ for all \hat{x} no matter which coordinates we choose.

□ Therefore, it is difficult to pinpoint in the matrix function $g_{\mu\nu}(x)$ the curvature degrees of freedom.

□ And: Even the entries $R^{\mu}_{\nu\sigma\tau}$ of the curvature tensor are coordinate system dependent.



Type here to search

4:14 PM
ENG
2020-02-11

is nontrivial just because of an unlucky choice of coordinates.

2. Spacetime is curved, i.e., we can not make $g_{\mu\nu}(x)$ take the form $\tilde{g}_{\mu\nu}(\hat{x}) = g_{\mu\nu}$ for all \hat{x} no matter which coordinates we choose.

□ Therefore, it is difficult to pinpoint in the matrix function $g_{\mu\nu}(x)$ the curvature degrees of freedom.

□ And: Even the entries $R^{\mu}_{\nu\sigma\tau}$ of the curvature tensor are coordinate system dependent.

Strategy:

□ Use the degrees of freedom of curvature to build a scalar and therefore coordinate system independent function $S_{\text{grav}}[g]$

□ Then, use this function as the action for gravity.

□ The equations of motion for gravity should follow from the action principle (and they do).

→ Need to define a scalar that encodes curvature!
We begin by going from a 4-tensor to a 2-tensor:

→ Need to define a scalar that encodes curvature!

We begin by going from a 4-tensor to a 2-tensor:

Definition: The "Ricci tensor:

$$R_{\mu\nu}(x) := R^i{}_{\mu i\nu}(x)$$

Recall: $\sum_{i=0}^3$ is implied

Note: Other index contractions would vanish because of antisymmetries of $R^i{}_{\nu\mu\varepsilon}(x)$ that are implied by the definition of $R^i{}_{\nu\mu\varepsilon}(x)$.

D...



$$R_{\mu\nu}(x) := R^i{}_{\mu i \nu}(x)$$

Recall: $\sum_{i=0}^3$ is implied

Note: Other index contractions would vanish because of antisymmetries of $R^i{}_{\nu\eta\varepsilon}(x)$ that are implied by the definition of $R^i{}_{\nu\eta\varepsilon}(x)$.

Remark:

$R_{\mu\nu}(x)$ carries strictly less information than the full Riemann curvature tensor:

- * If $R_{\mu\nu}(x)=0$ it is still possible that $R^i{}_{\nu\eta\varepsilon}(x) \neq 0$!
- * This happens to be the case, e.g., for gravitational waves.

Definition: The "curvature scalar" (or "Ricci scalar")

$$R(x) := g^{\mu\nu}(x) R_{\mu\nu}(x)$$

Other curvature scalars:

* The simplest scalar that can be formed from the metric alone is $g^{\mu\nu}(x) g_{\mu\nu}(x) = 4$.

* The next simplest scalar that can be formed is the Ricci scalar $R(x)$.

* All other scalars made out of g only are composed of higher powers of the Riemann tensor $R^\nu_{\mu\alpha\beta}(x)$:

$$R_{\mu\nu} R^{\mu\nu}, \quad R..R..R.. \text{ etc.}$$



The gravitational action

□ A priori, the full action now reads:

$$S'_{\text{tot}}[g, \phi, \psi_i] = S'_{\text{KG}} + S'_{\text{others}} + S'_{\text{grav}}$$

↑ other "matter" fields for e-, quarks, photons etc
↓ Klein-Gordon action

with: $S'_{\text{grav}}[g] := \int \left(c_0 + c_1 R(x) + c_2 R_{\mu\nu}(x) R^{\mu\nu}(x) + c_3 R^{\mu\nu} R_{\mu\nu} + \dots \right) \sqrt{|g|} d^4x$

□ Comparison with experiment shows evidence only for the first two terms:

$$S'_{\text{grav}}[g] = -\frac{1}{16\pi G} \int (2\Lambda + R(x)) \sqrt{|g(x)|} d^4x$$

↑ "Einstein action" ↑ Newton's constant ↓ "Cosmological constant"

Remark: D. Lovelock here determined all generalizations to higher terms and
at Applied Math at UW
 higher dimensions that still possess 2nd order initial value problems.

The equations of motion

The action principle is to require that the action be extremal with respect to all degrees of freedom:

$$A) \frac{\delta S_{\text{tot}}}{\delta \ell_i(x)} = 0 \quad B) \frac{\delta S'_{\text{tot}}}{\delta g_{\mu\nu}(x)} = 0 \quad C) \frac{\delta S_{\text{tot}}}{\delta \phi(x)} = 0$$

A) Require: $\frac{\delta S'_{\text{tot}}[g, \ell, \phi]}{\delta \ell_i(x)} = 0$

This yields the general relativistically covariant field equations for all "other" fields. (We will ignore the $\ell_i(x)$ for now.)

Quantization: Legendre transform $\rightarrow H(\ell_i, \pi_\ell) \rightarrow$ impose $\{\ell_i, \pi_\ell\} = i\delta_{i\ell}$.

B) Require: $\frac{\delta}{\delta g_{\mu\nu}(x)} S_{\text{eff}}[\phi, e, g] = 0$

This yields the equation of motion for the dynamics of curvature, i.e., the Einstein equation:

(See exercise
in Mukhanov's
text)

$$R_{\mu\nu}(x) - \frac{1}{2}g_{\mu\nu}(x)R(x) + \Lambda g_{\mu\nu}(x) = +8\pi G T_{\mu\nu}(x)$$

$$\sim \frac{\delta S'_{\text{grav}}}{\delta g_{\mu\nu}(x)}$$

$$\sim \frac{\delta(S'_{\text{curv}} + S'_{\text{Kc}})}{\delta g_{\mu\nu}(x)}$$

* Here, $T_{\mu\nu}(x)$ is the "Energy Momentum Tensor".
Neglecting the contribution by the $e_i(x)$, one obtains:

$$T_{\mu\nu}^{(\text{K.G.})}(x) = \phi_{,\mu}(x)\phi_{,\nu}(x) - g_{\mu\nu}(x) \left(\frac{1}{2}g^{\alpha\beta}(x)\phi_{,\alpha}(x)\phi_{,\beta}(x) - V(\phi(x)) \right)$$

mass term $m^2\phi^2$ included

* Quantization: To quantize the Einstein equation is difficult for many reasons:

- For example, it is difficult to separate the curvature degrees of freedom from mere artifacts of the choice of the coordinate system.
- Also, the Einstein equation is highly nonlinear.
- So far, all attempts have run into severe difficulties, even perturbative approaches.

→ This course: 1.) We will have initially consider known classical solutions $g_{\mu\nu}(x)$ and quantize only $\phi(x)$.
2.) Then, we will quantize linear perturbations of the metric.

c) Require: $\frac{\delta}{\delta \phi(x)} S_{\text{ext}}[g, \phi] = 0$

□ Since ϕ occurs only in S_{KG} we have, equivalently:

$$\frac{\delta S_{\text{KG}}}{\delta \phi(x)} = 0$$

□ Recall S_{KG} :

$$S_{\text{KG}}[\phi] = \frac{1}{2} \int_{\mathbb{R}^4} \left(g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2 - \lambda \phi^4 \right) \sqrt{|g|} d^4x$$

Example of a potential

□ Apply the Euler Lagrange equations:

$$\frac{\delta S_{\text{KG}}[\phi, \{\phi_{,\mu}\}, g]}{\delta \phi(x,t)} = \partial_\mu \frac{\delta S_{\text{KG}}[\phi, \{\phi_{,\mu}\}, g]}{\delta (\phi_{,\mu}(x,t))}$$



Type here to search

4:39 PM
ENG
2020-02-11

$\partial \psi(x)$

Since ϕ occurs only in S_{KG} we have, equivalently:

$$\frac{\delta S_{KG}}{\delta \phi(x)} = 0$$

Recall S_{KG} :

$$S_{KG}[\phi] = \frac{1}{2} \int_{\mathbb{R}^4} \left(g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2 - \frac{1}{2} \phi^4 \right) \sqrt{|g|} d^4x$$

Example of a potential

Apply the Euler Lagrange equations:

$$\frac{\delta S_{KG}[\phi, \{\phi_{,\mu}\}, g]}{\delta \phi(x,t)} = \partial_\mu \frac{\delta S_{KG}[\phi, \{\phi_{,\mu}\}, g]}{\delta (\phi_{,\mu}(x,t))}$$

□ \Rightarrow Klein Gordon equation in general relativity:

$$\left(-\frac{1}{2} m^2 \partial_\mu \phi(x) - \frac{1}{2} \lambda \phi^3(x) \right) \sqrt{g(x)} = \partial_\mu \left(\frac{1}{2} g^{\mu\nu}(x) \partial_\nu \phi(x) \sqrt{g(x)} \right)$$

Thus:

$$\frac{1}{\sqrt{g(x)}} \partial_\mu \left(g^{\mu\nu}(x) \sqrt{g(x)} \partial_\nu \phi(x) \right) + m^2 \phi(x) + \lambda \phi^3(x) = 0$$

□ Definition: The "d'Alembert operator", □

$$\square := \frac{1}{\sqrt{g(x)}} \partial_\mu g^{\mu\nu}(x) \sqrt{g(x)} \partial_\nu$$

Thus:

$$\square \phi(x) + m^2 \phi(x) + \lambda \phi^3(x) = 0$$

Next: Step 3 in

SR, 1st Q
Hamiltonian
formalism

step 1
Legendre transform
(equivalence)

SR, 1st Q.
Lagrangian
formalism

step 2
allow
curvature

GR, 1st Q
Hamiltonian
formalism

step 3
Legendre transform
(equivalence)

GR, 1st Q
Lagrangian
formalism

(PI)

GR, 2nd Q
Hamiltonian
formalism

Dyson-Schwinger eqns are same
(equivalence)

GR, 2nd Q
Lagrangian formalism
(Path integral of QFT)



Type here to search



4:42 PM
ENG
2020-02-11

Comment on step (PI): 2nd quantization with path integral

- Assume a fixed spacetime is chosen and we are given its metric $g_{\mu\nu}(x)$ in some arbitrary coordinate system.
- Then, for each field $\phi(\vec{x}, t)$ we can calculate its action $S_{\text{act}}[\phi, g]$:

$$S_{\text{act}}[\phi, g] = \frac{1}{2} \int_{\text{all}} \left(g^{\mu\nu} \partial_\mu \phi_{,\nu} - m^2 \phi^2 - \lambda \phi^4 \right) \sqrt{|g|} d^4x$$

- Following Feynman, we obtain "probability amplitudes":

$$\text{prob. ampl. } [\phi] := e^{\frac{i}{\hbar} S_{\text{act}}[\phi, g]}$$

□ Consider, e.g., the vacuum expectation value of $\phi(\vec{r}, t) \phi(\vec{r}', t)$, i.e., the correlation function of field amplitudes:

$$G(\vec{r}, t, \vec{r}', t) := \langle 0 | \hat{\phi}(\vec{r}, t) \hat{\phi}(\vec{r}', t) | 0 \rangle$$

□ We will later see how to calculate it using commutation relations etc.

□ With Feynman we also get it from the path integral:

Advantages:

- 1) Quick derivation of Feynman rules
- 2) Manifestly covariant.

Problems:

- 1) Ill defined due to uncountable number of integrations and related divergences.
- 2) Even when these are temporarily regularized, the identification of the measure still is ambiguous.
→ This issue is better handled in canonical formalism.

$$G(\vec{r}, t, \vec{r}', t) = N \int_{\text{all } \phi} \phi(\vec{r}, t) \phi(\vec{r}', t) e^{\frac{i}{\hbar} S_{\text{inv}}[\phi, g]} D[\phi]$$

↑ "Path Integral" over a function space



Type here to search

4:45 PM
ENG
2020-02-11

Recall strategy:

SR, 1stQ
Hamiltonian
formalism

GR, 1stQ
Hamiltonian
formalism

GR, 2ndQ
Hamiltonian
formalism

step 1
Legendre transform
(equivalence)

step 3
Legendre transform
(equivalence)

Dyson Schuring eqns are same
(equivalence)

SR, 1stQ.
Lagrangian
formalism

step 2
allow
curvature
GR, 1stQ
Lagrangian
formalism

GR, 2ndQ
Lagrangian formalism
(Path integral of QFT)

we are here

Recall:

General relativistic covariant Klein-Gordon theory in the Lagrangian formulation (neglecting the potential):

$$S_{KG} = \frac{1}{2} \int_{\mathbb{R}^4} \left(g^{\mu\nu}(x) \phi_{,\mu}(x) \phi_{,\nu}(x) - m^2 |\phi(x)|^2 \right) \sqrt{|g|} d^4x$$

we assume that the coordinate system is such, for simplicity.

It yields, via $\frac{\delta S_{KG}}{\delta \phi} = 0$ the Klein-Gordon eqn:

$$\frac{1}{\sqrt{|g(x)|}} \frac{\partial}{\partial x^\mu} \left(g^{\mu\nu}(x) \sqrt{|g(x)|} \phi_{,\nu}(x) \right) + m^2 \phi(x) = 0 \quad (KG)$$



* We read off the Lagrangian:

$$L_{\text{kg}}(t) = \frac{1}{2} \int_{\mathbb{R}^3} \left(g^{rr}(x,t) \dot{\phi}_{rr}(x,t) \dot{\phi}_{rr}(x,t) - m^2 \dot{\phi}^2(x,t) \right) \sqrt{g} d^3x$$

Step 3: Legendre transform back to the Hamiltonian form

* The transform:

$$H(\phi, \pi, t) \quad \xleftarrow[\text{Legendre transform}]{\pi(x,t) := \frac{\delta L}{\delta \dot{\phi}_{rr}(x,t)}} \quad L(\phi, \dot{\phi}_{rr}, t)$$

* Thus, the canonically conjugate field $\pi(x,t)$ reads:

$$\pi(x,t) = \frac{\delta L}{\delta \dot{\phi}_{rr}(x,t)} = \sqrt{|g(x,t)|} g^{rr}(x,t) \dot{\phi}_{rr}(x,t)$$

π the transform:

$$H(\phi, \pi, t) \xleftarrow[\text{Legendre transform}]{} \pi(x, t) := \frac{\delta L}{\delta \dot{\phi}_o(x, t)} \quad L(\phi, \dot{\phi}_o, t)$$

* Thus, the canonically conjugate field $\pi(x, t)$ reads:

$$\pi(x, t) = \frac{\delta L}{\delta \dot{\phi}_o(x, t)} = \sqrt{|g(x, t)|} g^{oo}(x, t) \dot{\phi}_{,o}(x, t)$$

* Explicitly:

$$\pi(x, t) = \sqrt{|g|} g^{oo} \dot{\phi}_{,o} + \sum_{i=1}^3 \sqrt{|g|} g^{oi} \dot{\phi}_{,i}$$

* Explicitly:

$$\pi(x,t) = \sqrt{|g|} g^{00} \phi_{,0} + \sum_{i=1}^3 \sqrt{|g|} g^{0i} \phi_{,i}$$

* Thus, we can also express $\phi_{,0}(x,t)$ in terms of $\phi(x,t)$ and $\pi(x,t)$ (as will be necessary after the Legendre transform):

$$\phi_{,0}(x,t) = \frac{\pi(x,t)}{g^{00}\sqrt{|g|}} - \sum_{i=1}^3 \frac{g^{0i}}{g^{00}} \phi_{,i}(x,t) \quad (\text{V})$$

↙ (of course, g^{00}
 depends on x, t too)

* The Hamiltonian:

$$H(\phi, \pi) = \int_{\mathbb{R}^3} \pi(x,t) \phi_{,0}(x,t) d^3x - \frac{1}{2} \int_{\mathbb{R}^3} \left(g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2 \right) \sqrt{|g|} d^3x$$

↑ Why do I need a



Type here to search

4:56 PM
ENG
2020-02-11

$$\pi(x,t) = \sqrt{|g|} g^{00} \phi_{,0} + \sum_{i=1}^3 \sqrt{|g|} g^{0i} \phi_{,i}$$

* Thus, we can also express $\phi_{,0}(x,t)$ in terms of $\phi(x,t)$ and $\pi(x,t)$ (as will be necessary after the Legendre transform):

$$\phi_{,0}(x,t) = \frac{\pi(x,t)}{g^{00}\sqrt{|g|}} - \sum_{i=1}^3 \frac{g^{0i}}{g^{00}} \phi_{,i}(x,t) \quad (\text{V})$$

↙ (of course, g^{00}
depends on x, t too)

* The Hamiltonian:

$$H(\phi, \pi) = \int_{\mathbb{R}^3} \pi(x,t) \phi_{,0}(x,t) d^3x - \frac{1}{2} \int_{\mathbb{R}^3} \left(g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2 \right) \sqrt{|g|} d^3x$$

Why don't we need a
factor of $\sqrt{|g|}$ for covariance
here? Because π has it built in!


$= L(\phi, \phi_{,0}(\phi, \pi), t)$

→ Exercise: Calculate $H(\phi, \pi, t)$ and simplify the expression as far as possible.

* The equations of motion:

We know from the general properties of the Legendre transform that the equations of motion now take the form:

$$\frac{d}{dt} \phi(x,t) = \frac{\delta H(\phi, \pi, t)}{\delta \pi(x,t)}, \quad \frac{d}{dt} \pi(x,t) = -\frac{\delta H(\phi, \pi, t)}{\delta \phi(x,t)}$$



$$\frac{d}{dt} \phi(x,t) = \frac{\delta H(\phi, \pi, t)}{\delta \pi(x,t)}, \quad \frac{d}{dt} \pi(x,t) = -\frac{\delta H(\phi, \pi, t)}{\delta \phi(x,t)}$$

* Exercise: Verify that these eqns are equivalent to (KG).

We are now ready to 2nd quantize:

SR, 1stQ
Hamiltonian
formalism

step 1
Legendre transform
(equivalence)

SR, 1stQ.
Lagrangian
formalism

GR, 1stQ
Hamiltonian
formalism

step 3
Legendre transform
(equivalence)

GR, 1stQ
Lagrangian
formalism

step 2
allow curvature




Type here to search



4:59 PM 2020-02-11

* Exercise: Verify that these eqns are equivalent to (KG).

We are now ready to 2nd quantize:



$$\begin{aligned}
 \int e^{i \int_{\text{tot}}' D\phi} &= \int e^{i \int_{\text{quadr.}}' + i \int_{\text{inter.}}'} D\phi \quad \left. \int x^n e^{-x^2 + jx} dx \right|_{j=0} \\
 &= \int e^{i \int_{\text{inter.}}(\phi) + i \int_{\text{quadr.}}(\phi)} D\phi \quad \text{IR} \\
 &= e^{i \int_{\text{inter.}}\left(\frac{i \delta}{8j}\right)} \int e^{i \int_{\text{quadr.}}(\phi) + i \int j\phi dx} D[\phi] \quad \left. \int e^{-x^2 + jx} dx \right|_{j=0} \\
 &\quad \text{Double } \left. \int e^{i \int j Q_j dx} \right|_{j=0} \quad \text{can do!} \sim e^{i \pi j^2}
 \end{aligned}$$

$$= \int_a^b f\left(\frac{d}{d\lambda}\right) e^{\lambda x} dx \Big|_{\lambda=0}$$

$$= f\left(\frac{d}{d\lambda}\right) \int_a^b e^{\lambda x} dx \Big|_{\lambda=0}$$

$\Rightarrow \int_a^b f(x) dx = f\left(\frac{d}{d\lambda}\right) \frac{-e^{\lambda x}}{\lambda} \Big|_{\lambda=0}$

Similarly: $\int_R f(x) dx = \lim_{\varepsilon \rightarrow 0} \frac{f(\partial_\varepsilon) + \int_{\partial_\varepsilon}^x f(x) dx}{\varepsilon}$

$$\int_{-\infty}^{\infty} f(x) dx$$



$$\begin{aligned} \int_a^b f(x) dx &= \int_a^b f(x) e^{\lambda x} dx \Big|_{\lambda=0} \\ &= \int_a^b f\left(\frac{d}{d\lambda}\right) e^{\lambda x} dx \Big|_{\lambda=0} \\ &= f\left(\frac{d}{d\lambda}\right) \int_a^b e^{\lambda x} dx \Big|_{\lambda=0} \end{aligned}$$

$$\left. \int_{-\infty}^{\infty} f(x) dx = 2\pi \int (-i\partial_x) \delta(x) \right|_{x=0} \quad \left. \int_a^b f(x) dx = \int_a^b f(x) e^{\lambda x} dx \right|_{\lambda=0}$$

$$= \int_a^b f\left(\frac{d}{d\lambda}\right) e^{\lambda x} dx \quad \left. \right|_{\lambda=0}$$

$$= f\left(\frac{d}{d\lambda}\right) \left. \int_a^b e^{\lambda x} dx \right|_{\lambda=0}$$

Similarly:

$$\int_{\mathcal{R}} f(x) dx = \lim_{\varepsilon \rightarrow 0} \frac{f(\partial_\varepsilon) + f(-\partial_\varepsilon)}{\varepsilon}$$