Title: A Symmetry of Cosmological Observables: Looking at Hubble through the Mirror

Speakers: Francis-Yan Cyr-Racine

Series: Cosmology & Gravitation

Date: April 19, 2022 - 11:00 AM

URL: https://pirsa.org/22040121

Abstract: Many key cosmological observables have a built-in symmetry under rescaling of all important length scales in the problem. This scaling symmetry can be used to make partial progress toward a complete resolution of the Hubble tension. To exploit the symmetry while respecting observational constraints, we are naturally led to a "mirror world" dark sector. A successful implementation of this scaling symmetry requires a means of increasing the cosmic photon scattering rate that respects observational bounds on the primordial helium abundance. We discuss different ideas that could in principle achieve this rescaling, and discuss their advantages and drawbacks. We finally present some general observations about the fundamental nature of the "Hubble tension".

Zoom Link: https://pitp.zoom.us/j/99678466262?pwd=VTY4cWNFUERhU08vYi9lT09MZjJkdz09

Pirsa: 22040121 Page 1/40

A Symmetry of Cosmological Observables: Looking at Hubble through the Mirror Sector



Perimeter Institute

Francis-Yan Cyr-Racine


Department of Physics and Astronomy, University of New Mexico

With Fei Ge (UC Davis), Lloyd Knox (UC Davis), and Kylar Greene (UNM)



Pirsa: 22040121 Page 2/40

Hubble rate: The Landscape



Francis-Yan Cyr-Raci...

Freedman (2021)


Francis-Yan Cyr-Racine - UNM

4/19/22

Pirsa: 22040121 Page 3/40

Hubble tension: Latest SH0ES results





Riess et al., arXi:2112.04510

- Now SH0ES nearly 5-sigma away from Planck CMB!
- The seriousness of the tension is now difficult to dismiss.
- Exciting times for cosmology!

Francis-Yan Cyr-Racine - UNM

4/19/22


3

Pirsa: 22040121 Page 4/40


Why you should take this seriously



• Among other important improvements, the doubling of the calibrator supernova host sample is a major step forward.







Riess et al., arXi:2112.04510

Francis-Yan Cyr-Racine - UNM

4/19/22

4

Pirsa: 22040121 Page 5/40

Where does this leave us?

Francis-Yan Cyr-Raci...

- According to this scale, we are in a crisis
 - $< 1\sigma$: Consistency
 - $> 2\sigma$: Curiosity
 - $> 3\sigma$: Tension/discrepancy
 - > 4σ : Problem
 - $\gtrsim 5\sigma$: Crisis





Francis-Yan Cyr-Racine - UNM


4/19/22

Theorists to the rescue!



In this crisis, our job as theorists is to identify what properties a successful solutions might have.





Bottom line:

We have yet to identify a complete solution that is palatable to both cosmologists and particle physicists but have found important clues about what a successful model would look like.

(quote from KITP Conference July 2019)

Francis-Yan Cyr-Racine - UNM

4/19/22

6




Pirsa: 22040121 Page 7/40

Theorists to the rescue!

Niayesh Afshordi

In this crisis, our job as theorists is to identify what properties a successful solutions might have.





Bottom line:

We have yet to identify a complete solution that is palatable to both cosmologists and particle physicists but have found important clues about what a successful model would look like.

(quote from KITP Conference July 2019)

Francis-Yan Cyr-Racine - UNM


4/19/22

6

Pirsa: 22040121 Page 8/40

This talk: Why is it so hard to get a large Hubble constant from the CMB + BAO?





• Lots of ideas out here! Why are they struggling to get a large value of the Hubble rate?

Schöneberg et al., arXi:2107.10291


Francis-Yan Cyr-Racine - UNM

4/19/22

Late solution: Could we just do that?

See e.g. Di Valentino et al. (2021






Francis-Yan Cyr-Racine - UNM


4/19/22

No!! Any model must fit the **distances** to type Ia supernovae



• Just increasing the Hubble rate at very late times **does not** address the root cause of the tension.





Benevento et al. (2020)

Greene and Cyr-Racine, arXiv:2112.11567

Francis-Yan Cyr-Racine - UNM


4/19/22

Pirsa: 22040121 Page 11/40

No!! Any model must fit the **distances** to type Ia supernovae



• Not properly taking distances into account can lead to the detection of spurious physics!



• >3 sigma detection of a late transition in the abundance of dark energy (bogus!)

Greene and Cyr-Racine, arXiv:2112.11567

Francis-Yan Cyr-Racine - UNM

4/19/22

10

Pirsa: 22040121 Page 12/40

Important Lesson




If you are working on a late-time solution, please make sure that your model fit the actual distances to Type Ia supernovae.

Francis-Yan Cyr-Racine - UNM

4/19/22

11

Pirsa: 22040121 Page 13/40



Pirsa: 22040121 Page 14/40


Baryon Acoustic Oscillations (BAO)



BAO primarily measures 2 "processed" versions of the baryon-photon sound horizon.

Line of sight: $H(z)r_{
m s}$

Transverse: $r_{
m s}/D_A(z)$



If sound horizon is known, then we can use BAO to infer Hubble rate.

Francis-Yan Cyr-Racine - UNM


4/19/22

13

Pirsa: 22040121 Page 15/40

CMB and BAO measurements of the expansion rate are all anchored on the sound horizon





Important:

The baryon-photon sound horizon, r_s , is something we compute theoretically, assuming we understand atomic physics and the energy content of the Universe.

Time of baryon decoupling

$$r_s = \int_0^{t_{\rm d}} c_{\rm s} dt / a = \int_0^{a_{\rm d}} c_{\rm s} \frac{da}{a^2 H(a)}$$

Francis-Yan Cyr-Racine - UNM


4/19/22

14

Pirsa: 22040121 Page 16/40

Early solution: Cartoon





Francis-Yan Cyr-Racine - UNM

4/19/22

15

Pirsa: 22040121 Page 17/40

Question




Can we change the cosmological expansion history and leave the CMB/BAO *exactly* invariant?

Francis-Yan Cyr-Racine - UNM

4/19/22

16

Pirsa: 22040121 Page 18/40



Pirsa: 22040121 Page 19/40

However, the CMB as a whole is not invariant under $H \rightarrow fH$



- Why? Photon-baryon equations of motion depend on two other dimensionful quantities: wavenumber *k* and the baryon-photon scattering rate.
- For example, let's look at the continuity equation for photons:

(Use scale factor "a" as time variable.)


Francis-Yan Cyr-Racine - UNM

4/19/22

However, the CMB as a whole is not invariant under $H \rightarrow fH$



- Why? Photon-baryon equations of motion depend on two other dimensionful quantities: wavenumber *k* and the baryon-photon scattering rate.
- For example, let's look at the continuity equation for photons:



Francis-Yan Cyr-Racine - UNM

4/19/22

However, the CMB as a whole is not invariant under $H \rightarrow fH$



- Why? Photon-baryon equations of motion depend on two other dimensionful quantities: wavenumber *k* and the baryon-photon scattering rate.
- For example, let's look at the Euler equation for photons:

Photon heat flux Photon quadrupole (Use scale factor "a" as time variable.) $a^2H\frac{\partial F_{\gamma 1}}{\partial a}=\frac{k}{3}(F_{\gamma 0}-2F_{\gamma 2})+\frac{4k}{3}\psi+\dot{\kappa}(\frac{4}{3}v_{\rm b}-F_{\gamma 1}),$ Photon temperature monopole perturbation Gravitational driving term Photon-baryon scattering rate

Francis-Yan Cyr-Racine - UNM


4/19/22



Symmetry: Basic geometry and the dimensional analysis



- Dimensionless observables seen in projection on the sky have an intrinsic scale invariance.
- By dimensional analysis,
 ODEs for the evolution of
 dimensionless quantities can
 only depend on
 dimensionless ratios.



Francis-Yan Cyr-Racine - UNM

4/19/22


20

Pirsa: 22040121 Page 23/40

Symmetry: Basic geometry and the dimensional analysis



- Dimensionless observables seen in projection on the sky have an intrinsic scale invariance.
- By dimensional analysis,
 ODEs for the evolution of
 dimensionless quantities can
 only depend on
 dimensionless ratios.



Nothing special about cosmology here!

Francis-Yan Cyr-Racine - UNM 4/19/22 20

Pirsa: 22040121 Page 24/40

But what if **all** length scales are uniformly rescaled?



• Instead of just rescaling *H*, do the transformation:

$$H \to fH, k \to fk, \dot{\kappa} \to f\dot{\kappa}.$$

• By dimensional analysis, all factors of f cancel out in the equations of motion (EOM).

This leaves the photon-baryon (and dark matter and massless neutrinos) EOM invariant.

Francis-Yan Cyr-Racine - UNM

4/19/22

Special feature of our Universe: Initial conditions



• We happen to live in a Universe in which the initial scalar fluctuations have no intrinsic scale.

$$P(k) = A_{\rm s}(k/k_{\rm p})^{n_{\rm s}-1}$$

- Since $n_s < 1$, the different Fourier modes have slightly different primordial amplitudes.
- Thus, the transformation $k \to fk$ will modify the amplitude of fluctuations (CMB, $P_{\rm m}(k)$, etc.)
- However, since power laws have no scale, this can be corrected with a trivial rescaling:

Zahn and Zaldarriaga (2003)

$$A_{
m s}
ightarrow A_{
m s}/f^{n_{
m s}-1}$$

Francis-Yan Cyr-Racine - UNM

4/19/22

22

Pirsa: 22040121 Page 26/40



The scaling "recipe"

1. Increase Hubble rate at all times by scaling up every energy density:

$$G
ho_i o f^2G
ho_i$$

2. Scale up the photon scattering rate $\dot{k} = a n_e \sigma_T$ according to:

$$\sigma_{\rm T} n_e(a) \to f \sigma_{\rm T} n_e(a)$$

3. Adjust the initial amplitude of scalar fluctuations according to

$$A_{
m s} o A_{
m s}/f^{n_{
m s}-1}$$

Cyr-Racine, Ge, Knox, arXiv: 2107:13000, PRL accepted.

Francis-Yan Cyr-Racine - UNM

4/19/22








This works



This really leaves the CMB temp/pol invariant (fixing recombination history here)








$$H_0 = 67.5, 74.3, 81 \text{km/s/Mpc}$$

Francis-Yan Cyr-Racine - UNM

4/19/22




Pirsa: 22040121 Page 29/40

Getting around COBE: Mirror World



- We can't easily increase the densities of photons/baryons
- So instead add mirror "dark" particles!



Stranger Things

Chacko et al. (2005, a,b,c), Craig & Howe (2014), Craig et al. (2015), Farina (2015), Barbieri et al. (2016), Chacko et al. (2017), Csaki et al. (2017), Hochberg et al. (2017), Harigaya et al. (2017), Ibe et al. (2019), Terning et al. (2019), Curtin & Gryba (2021), Blinov et al. (2021) and many more

Francis-Yan Cyr-Racine - UNM

4/19/22

26

Pirsa: 22040121 Page 30/40

Adjusting the photon scattering rate



• The mirror world ingredients ensure that we "effectively" implement the rescaling

$$\sqrt{G
ho_i(a)}
ightarrow f\sqrt{G
ho_i(a)}, \quad \blacksquare
ightarrow H
ightarrow fH$$

while leaving the perturbation evolution invariant.

• It does not however implement the necessary scaling

$$\sigma_{\rm T} n_e(a) \to f \sigma_{\rm T} n_e(a)$$

• Since $n_e \propto (1 - Y_P)$, one can implement this scaling by adjusting the helium fraction according to

$$(1 - Y_{\rm P}) \to f(1 - Y_{\rm P})$$


Unlikely to be the right way of doing this!

Francis-Yan Cyr-Racine - UNM

4/19/22








- With a fixed recombination history, we can fit any value of the Hubble constant to CMB + BAO data.
- This is just a numerical realization of the symmetry we already know exists.

Francis-Yan Cyr-Racine - UNM

4/19/22







• Symmetry-breaking effects from the self-consistently calculated recombination history are very mild!

Francis-Yan Cyr-Racine - UNM


4/19/22

29

Pirsa: 22040121 Page 33/40

Second Test: Compatibility with the cepheid-calibrated distance ladder





- In principle, the symmetry allows us to completely eliminate the Hubble tension between CMB + BAO and the local distance ladder (R21 here).
- However, we are doing the scaling by fiat here.

Francis-Yan Cyr-Racine - UNM

4/19/22





31

- Three key ingredients are necessary to turn this general scaling transformation into an actual model
 - A mirror dark sector that (nearly) mimic the SM (see e.g. Blinov et al., arXiv:2108.11386)
 - A means to rescale the photon scattering rate other than helium (see e.g. Sekiguchi and Takahashi, arXiv:2007.03381; Burgess et al. arXiv:2111.07286).
 - A means to ensure consistency with helium and deuterium abundance measurements.

Francis-Yan Cyr-Racine - UNM 4/19/22

Pirsa: 22040121 Page 35/40





- Three key ingredients are necessary to turn this general scaling transformation into an actual model
 - A mirror dark sector that (nearly) mimic the SM (see e.g. Blinov et al., arXiv:2108.11386)
 - A means to rescale the photon scattering rate other than helium (see e.g. Sekiguchi and Takahashi, arXiv:2007.03381; Burgess et al. arXiv:2111.07286).
 - A means to ensure consistency with helium and deuterium abundance measurements.

These provide clear model-building targets!

Francis-Yan Cyr-Racine - UNM

4/19/22


31

Pirsa: 22040121 Page 36/40

Possible implication: Mirror Sector Freedom

Francis-Yan Cyr-Raci...

At face value, the direct Hubble measurements predict ~3% in atomic dark matter, and a dark photon bath with a neutrino-like temperature.



Cyr-Racine, Ge, Knox, arXiv: 2107:13000, PRL accepted

4/19/22


32

Francis-Yan Cyr-Racine - UNM ← △ ్ ⊶

Pirsa: 22040121 Page 37/40

The scaling symmetry helps understand constraints on other cosmological models





Schöneberg et al., arXiv:2107.10291

Francis-Yan Cyr-Racine - UNM

4/19/22

33

Pirsa: 22040121 Page 38/40

Conclusions



- The Hubble tension is really about distance measurements.
- Early solutions must be able to fit distances to SNe Ia.
- We have identified a previously unnoticed symmetry of cosmological observables projected on the sky.
- If a complete model could be found, it would completely eliminate the Hubble tension.
- This symmetry helps us understand the nature of cosmological constraints.

Francis-Yan Cyr-Racine - UNM

4/19/22

34

Pirsa: 22040121 Page 39/40



35

Open Questions

- Can we achieve a higher photon scattering rate and have consistency with BBN and Y_P?
- Can we detect the 3% of atomic DM?
- Can a consistent mirror sector be built? (Yes!)
- Impact of nonlinear evolution?

Francis-Yan Cyr-Racine - UNM 4/19/22

Pirsa: 22040121 Page 40/40