

Title: Session 2 - Youssef Mousaaid


Speakers: Youssef Mousaaid

Collection: POSTDOC WELCOME 2022

Date: October 24, 2022 - 11:30 AM

URL: <https://pirsa.org/22100123>

2- Categories



Set

category

functions

functors

equations

isomorphisms

decat

K_0

Grothendieck


tr

trace functor

$$g \longrightarrow \mathcal{U}_q(\mathfrak{g}) \longrightarrow \hat{\mathcal{U}}_q(\mathfrak{g})$$

KLR defined 2-cat \mathcal{E}_g s.t. $K_0(\mathcal{E}_g) = \hat{\mathcal{U}}_q(\mathfrak{g})$

Notations (Crane-Frenkel)




$$L \simeq L' \implies Kh(L) \simeq Kh(L')$$

$$L \longrightarrow Kh(L) \longrightarrow [Kh(L)] \in K_0(\mathcal{E}_g)$$

$$\begin{array}{ccc} \parallel & & \parallel \\ \mathcal{J}_n(L) & \simeq & \mathbb{Z}[q, q^{-1}] \end{array}$$

Categorify
 Skein model of the
 torus



$g \rightarrow \mathcal{U}_q(\mathfrak{g}) \rightarrow \mathcal{U}_q(\mathfrak{g})$

KLR defined 2-cat \mathcal{E}_g s.t. $K_0(\mathcal{E}_g) = \mathcal{U}_q(\mathfrak{g})$

Motivations (Crane-Frenkel)

$\mathcal{I}_q(L) \xrightarrow{\text{WRT}} \mathbb{Z}^n$ invariant

K_h

$\mathcal{K}_h(L)$

$\mathcal{M}(L)$

$L \cong L' \rightarrow Kh(L) \cong Kh(L')$

$L \rightarrow Kh(L) \rightarrow [Kh(L)] \in K_0(\mathcal{E}_g)$

\parallel

$\mathbb{Z}^n(L) \cong \mathbb{Z}[q, q^{-1}]$

Categorify

Skein model of the

torus