Title: Analogue quasinormal mode oscillations of optical solitons

Speakers:

Collection: Quantum Simulators of Fundamental Physics

Date: June 08, 2023 - 9:00 AM

URL: https://pirsa.org/23060017

Analogue Quasinormal Mode Oscillations of Optical Solitons

Friedrich Koenig

P.I. Workshop Waterloo June 5-9

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

SUPA

QSimFP

EPSRC
Engineering and Physical Sciences Research Council

Contents

Motivation

- Introduction to Ringdown
- Example: *Black Hole Oscillations*

- Ringdown and Quasinormal modes
- Black hole Quasinormal modes
- Optical Soliton QNMs
- Soliton / Black Hole Analogy
- Ringdown in Optical Solitons
- Summary and Outlook

Jack Petty
St. Andrews

Christopher Burgess
St. Andrews

Pavlos Manousiadis
St. Andrews

Sam Patrick
King's College

Ruth Gregory
King's College

Theo Torres
Aix-Marseille

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Systems with Ringdown

Many perturbative systems relax in oscillatory way

church bells
leaky cavities
black holes
etc.

stringed instruments
electrical circuits with resistance
water vortex

❖ **Use in string theory, brane-world models, quantum gravity**

Resonator Ringdown Following an Initial Perturbation
Lalanne, P., Yan, W., Vynck, K., Sauvan, C., & Hugonin, J. P. (2018)



Helmholtz Resonator

Spectrum of light ring modes in a water vortex



S. Patrick et al., PRL **121**, 061101 (2018).
Th. Torres et al. PRL **125**, 011301 (2020).

R. A. Konoplya and A. Zhidenko, Rev. Mod. Phys. **83**, 793 (2011).
E. Berti et al., Class Quantum Grav. **26,** 163001 (2009).
C. Sauvan et al., Opt. Ex., **30**, 6846 (2022).

Sagittarius A*, the black hole at the center of the Milky Way galaxy. EHT Collaboration

Black Hole Oscillations

Perturbed black hole "rings down" like a church bell

The bell rings from its material parts
The black hole rings from the spacetime near its light-ring

Both have a *characteristic sound*



Black Hole Spacetime

$c = G = 1$

Schwarzschild $ds^2 = -\left(1 - \frac{r_s}{r}\right) dt^2 + \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 + r^2 d\Omega$ ***(Kerr, Reissner–Nordström, Kerr–Newman) for any asymptotic curvature***

Perturbation Field

e.g. scalar, electromagnetic, gravitational → ***emission "ringdown" waves***[5]

$$g_{\mu\nu} = g^0_{\mu\nu} + \delta g_{\mu\nu}$$

[5] LIGO, 2016

Supermassive Black Hole M87*

The Event Horizon Telescope Collaboration et al. ApJL, 875 L4, (2019)

Black Hole Oscillations

PERTURBED BLACK HOLE



LIGO DETECTOR



LIGO SIGNAL



LIGO Detection of a Black Hole Merger



Best-Fit Theoretical Waveform to LIGO data *PRL 116, 061102 (2016)*

"Hearing the shape of a drum"

Can we hear the shape of a black hole?

Normal Modes v Quasinormal Modes

NORMAL MODE



$$x(t) \sim \underbrace{e^{-i\omega t}}_{\text{oscillation}}$$



→ ***closed***
→ ***conservative***
→ ***Hermitian***

Examples

harmonic oscillators
quantum fields

QUASINORMAL MODE (QNM)



$$x(t) \sim \underbrace{e^{-\Gamma t}}_{\text{decay}} \underbrace{e^{-i\omega t}}_{\text{oscillation}}$$

$$\Omega = \omega - i\Gamma$$

→ ***open***
→ ***non-conservative***
→ ***non-Hermitian***

Examples

plasmonic resonators [1]
fluid vortices [2]
black holes [3]
optical solitons

[1] Lalanne, 2018
[2] Torres, 2018
[3] Berti, 2009

www.st-Andrews.ac.uk/~qoi

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Quasinormal Modes of Fields

LOCALISED SYSTEMS



System

→ ***localised system***
e.g. time-independent potential
cavity / resonator

→ ***imperfect store of energy***
QNMs are "leaky modes"

Quasinormal Modes

→ ***outgoing energy flux***
→ ***decaying in time***
→ ***growing in space***
→ ***unphysical***

QUASINORMAL MODE SHAPE



$$\psi \sim e^{-i\omega(t \pm x/v)}$$

[4] Kokkotas, Schmidt, L. Rev. Rel. 1999

Ringdown and Quasinormal Modes

Ringdown as relaxation of a perturbed system

$$\partial_t^2\psi(t,x) + (-\partial_x^2 + V(x))\psi(t,x) = 0$$

Initial conditions $\psi(0,x),\ \partial_t\psi(0,x)$

Solution with Greens function:

$$\psi(t,x) = \frac{1}{2\pi i}\int_{-\infty}^{\infty} e^{(a+i\Omega)t}\int_{-\infty}^{\infty} G(a+i\Omega,x,x')\,j(a+i\Omega,x')\quad dx'd\Omega$$

Greens function Initial data



Ching et al., 1995

→ ***QNMs manifest in linear response***

$$\psi(t,x) \sim \sum_{n=0} a_n e^{i\Omega_n t} u(\Omega_n,x) + \psi_{rest}(t,x)$$

QNMs

QNM expansion develops over finite spaces [4]



[4] Kokkotas, Schmidt, L. Rev. Rel. 1999

Effective Gravitational Potential



Perturbation Field

e.g. *Massless Scalar Field* $\frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}\,g^{\mu\nu})\partial_\nu\psi = 0$

Radial Wavefunction

$$\psi(t,r,\theta,\phi) = e^{-i\omega t}\frac{\Psi(r)}{r}Y_l(\theta,\phi)$$

1D Schrödinger Equation

$$\partial_{r_*}^2\Psi + (\omega^2 - V)\Psi = 0$$

Effective Gravitational Potential $V(r) = \left(1-\frac{r_s}{r}\right)\left[\frac{l(l+1)}{r^2}+\frac{r_s}{r^3}\right]$

(peaked around light-ring)

Tortoise co-ordinate $\frac{dr}{dr_*} = \left(1-\frac{r_s}{r}\right)$

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Mashoon: Pöschl-Teller approximation

$$\frac{d^2\Psi_s}{dr_*^2} + (\omega^2 - V_s)\Psi_s = 0 \qquad V_s = \left(1 - \frac{2M}{r}\right)\left[\frac{l(l+1)}{r^2} + (1-s^2)\frac{2M}{r^3}\right]$$



`Low lying' QNMs are dominated by the peak of the potential.



Exact solutions available for Pöschl-Teller potential [1]:

$$\omega_n = \pm\sqrt{V_0 - \alpha^2/4} - i\alpha(2n+1)/2$$

$$n = 0, 1, 2, \ldots$$

$$V_{\rm PT} = \frac{V_0}{\cosh^2\alpha(r_* - r_{*0})}$$



[1] H.-J. Blome and B. Mashhoon, Phys. Lett. **110A**, 231 (1984).

Black Hole QNM Spectrum

Pöschl-Teller Approximation[6]

$$V \approx V_{\mathrm{PT}} = V_0 \,\mathrm{sech}^2\, \alpha(r_* - r_{*0})$$



QNM Spectrum

$$\omega_n = \pm\sqrt{V_0 - \alpha^2/4} - i\alpha\left(n + \frac{1}{2}\right)$$

$n = 0, 1, 2, \ldots$



→ ***Schwarzschild***
→ ***SdS***
→ ***SAdS***
→ ***Nariai***

[6] Konoplya, 2011

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Optical Solitons

Pulse Propagation in Dispersive Kerr Media[7]

Optical Kerr effect

$$n(\omega, t) = n_0(\omega) + n_2 I(t)$$



Nonlinear Schrödinger Equation (higher-order dispersion)

$$\partial_z A - i[\beta(i\partial_t + \omega_s) - \beta(\omega_s)]A - i\gamma|A|^2 A = 0$$

$\beta(\omega) = \frac{n_{\text{eff}}(\omega)\,\omega}{c}$	propagation constant
ω_s	carrier frequency
γ	fibre nonlinear coefficient

Soliton Solution

$$A_s(z,t) = \sqrt{P_0}\ \text{sech}\left(\frac{t - z/v}{T_0}\right) e^{i\gamma P_0 z}$$



approximate solution

P_0	soliton pulse peak power
v	group velocity at soliton frequency
T_0	soliton temporal pulse width

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

St Andrews

[7] Agrawal, 2003

Optical analogue to QNMs?

Wave velocity in moving frame: optical pulse in Kerr medium

A pulse moves at velocity v_0 and locally modifies the refractive index (i.e. wave velocity) by the cross-Kerr effect.







University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Soliton as Optical Potential

Linear Perturbation Equation

$$\partial_z A_p - i[\beta(i\partial_t + \omega_s) - \beta(\omega_s)]A_p - 2i\gamma|A_s|^2 A_p - i\gamma A_s^2 A_p^* = 0$$

fast-oscillating

$$A = A_s + A_p \qquad (|A_p|^2 \ll P_0)$$

LABORATORY FRAME



Group-Velocity Matched Perturbation Equation

→ ***spectrally distinct*** ⟶ neglect $i\gamma A_s^2 A_p^*$

→ ***narrowband*** ⟶ $\beta(\omega) = \sum_{k=0}^{2} \frac{\beta_{ak}}{k!}(\omega - \omega_a)^k$

→ ***group velocity matched*** ⟶ $\beta_{a1} = \beta_{s1} = 1/v$

MOVING FRAME



$$\beta_{s1}\partial_\zeta a_p + \frac{i}{2}\frac{\beta_{a2}}{T_0^2}\partial_\tau^2 a_p - 2i\gamma|A_s|^2 a_p = 0$$

ω_a perturbation freq.

soliton intensity as potential

moving frame

$$\tau = \frac{t - z/v}{T_0} \qquad \zeta = z/v$$

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland
www.st-Andrews.ac.uk/~qoi

Soliton Quasinormal Modes

Soliton GVM Modes

$$a_p(\zeta,\tau) = u(\tau)e^{-i\Omega\zeta}$$

$$\partial_\tau^2 u + \left[-\frac{2|\beta_{s2}|\Omega}{\beta_{a2}} - \frac{4|\beta_{s2}|}{\beta_{a2}}\operatorname{sech}^2(\tau)\right]u = 0$$

QNM Solution

$$a_{p,n}(\zeta,\tau) = \underbrace{\cosh(\tau)^{n+1/2}}_{1} \times \underbrace{\exp\left(\operatorname{Im}(\Omega_n)\zeta\right)}_{2} \times \underbrace{\exp\left(-i\operatorname{Im}(\lambda)\log\cosh(\tau) - i\operatorname{Re}(\Omega_n)\zeta\right)}_{3} \times \underbrace{{}_2F_1[-n, 1+2\lambda-n; 1+\lambda-n; [1-\tanh(\tau)]/2]}_{4}$$

discrete spectrum
$n = 0, 1, 2, \ldots$

1. Divergence in τ ('space')
2. Decay in the ζ ('time')
3. Directs phase velocity
4. Order-n polynomial in $\tanh(\tau)$ controls parity of solution

$$\lambda = -\frac{1}{2} + i\sqrt{\frac{4|\beta_{s2}|}{\beta_{a2}} - \frac{1}{4}}$$

'SPATIAL' PROFILE



'TEMPORAL' PROFILE



University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Soliton QNM Spectrum

QNM Frequencies

$$\Omega_n = \frac{\beta_{a2}}{2|\beta_{s2}|}\left[\left(n+\frac{1}{2}\right)^2 - \left(\frac{4|\beta_{s2}|}{\beta_{a2}} - \frac{1}{4}\right)\right] - i\frac{\beta_{a2}}{|\beta_{s2}|}\left(n+\frac{1}{2}\right)\sqrt{\frac{4|\beta_{s2}|}{\beta_{a2}} - \frac{1}{4}}$$

discrete spectrum

$n = 0, 1, 2, \ldots$

→ ***QNM overtones fall along a parabola***

$$\mathrm{Im}(\Omega)^2 - \left(8 - \frac{\beta_{a2}}{2|\beta_{s2}|}\right)\mathrm{Re}(\Omega) + \frac{1}{4}\left(8 - \frac{\beta_{a2}}{2|\beta_{s2}|}\right)^2 = 0$$

→ ***Finitely many outgoing QNMs***

$$N_{\mathrm{out}} = \left\lfloor \frac{1}{2} + \sqrt{\frac{4|\beta_{s2}|}{\beta_{a2}} - \frac{1}{4}} \right\rfloor$$

→ ***Infinitely many ingoing QNMs***



University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Soliton QNM Representations II



$n = 2$	$n = 3$	$n = 4$

1. Higher modes diverge in τ more quickly.
2. Higher modes decay in ζ more quickly.
3. **Phase velocities turn around and then increase.**



soliton background



University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Optical Soliton / Black Hole QNM Analogy

Optical Soliton QNMs

→ ***new black hole analogue system***

Formally Equivalent Mode Equations

$$\partial_\tau^2 u + \left[-\frac{2|\beta_{s2}|\Omega}{\beta_{a2}} - \frac{4|\beta_{s2}|}{\beta_{a2}} \operatorname{sech}^2(\tau) \right] u = 0$$

GVM Soliton QNMs

$$\partial_\rho^2 \Psi + \left[\frac{\omega^2}{\alpha^2} - \frac{V_0}{\alpha^2} \operatorname{sech}^2(\rho) \right] \Psi = 0$$

Black Hole QNMs

→ ***identical spatial profiles***

- *identical field expansions*

$$\Omega \sim -\omega^2$$

→ ***different mode frequencies***

- *different temporal profiles*
- *different field evolution*

(dispersionless limit may solve this)

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Soliton QNMs: Dispersionless Limit

BLACK HOLE QNM SPECTRUM



$$\omega^2 \sim -\Omega$$

OPTICAL SOLITON QNM SPECTRUM



QNM frequencies become equivalent for small dispersion at the perturbation:

$$|\beta_{s2}| \gg \beta_{a2}$$

$$\omega_n = \pm\sqrt{V_0 - \frac{\alpha^2}{4}} - i\alpha\left(n + \frac{1}{2}\right) \qquad \Omega_n \approx -2 - i\sqrt{\frac{4\,\beta_{a2}}{|\beta_{s2}|}}\left(n + \frac{1}{2}\right)$$



MORE DISPERSION

LESS DISPERSION

University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Simulation: Ringdown in Optical Solitons

SCATTERING OF DISPERSIVE WAVE



RINGDOWN SIGNAL



RINGDOWN SPECTRUM



→ ***near-GVM pulse on soliton***

all pulse shapes excite QNMs

best: *super-gaussian and chirped*

$$V \equiv \frac{4|\beta_{s2}|}{\beta_{a2}}$$

→ ***ringdown in wake of transmission/reflection***

REAL PART



IMAGINARY PART



University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

QNM with evolving soliton

Rerturbation field (real part, soliton background removed)



RINGDOWN SIGNAL



University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland

Summary and Outlook

Summary

- Soliton acts as an optical potential
- NLS solitons exhibit “ringdown” described by QNMs
- New experimentally accessible black hole analogue
- First soliton QNM system & first with “ingoing” QNMs



Outlook

- Soliton ringdown to be observed experimentally
- Generalizations beyond GVM QNMs
- QNMs of other solitons (e.g. KdV, sine-Gordon)
- Quantum effects with QNMs



University of St Andrews
St Andrews, Fife, KY16 9AJ, Scotland