

Title: The Two Scales of New Physics in Higgs Couplings

Speakers: Gabriele Rigo

Series: Particle Physics

Date: October 27, 2023 - 1:00 PM

URL: <https://pirsa.org/23100112>

Abstract: Higgs coupling deviations from Standard Model predictions contain information about two scales of Nature: that of new physics responsible for the deviation, and the scale where new bosons must appear. The two can coincide, but they do not have to. The scale of new bosons can be calculated by going beyond an effective field theory description of the coupling deviation. I will discuss model-independent upper bounds on the scale of new bosons for deviations in Higgs to WW and ZZ couplings, and explain how any measured deviation at present or future colliders requires the existence of new bosons within experimental reach. This has potentially interesting implications for naturalness.

Zoom link <https://pitp.zoom.us/j/97203814123?pwd=U3F3N2xxTmJSQmxIQ0Rib3ZwN3Fldz09>

The Two Scales of New Physics in Higgs Couplings

[arXiv:2305.19325 [hep-ph]]

Gabriele Rigo (IPhT, Saclay)

Raffaele Tito D'Agnolo, Florian Nortier, Pablo Sesma




Particle Physics Seminar

27 October 2023




Basic Questions

Origin of mass?



Hierarchy problem

Composition of the Universe?



Dark matter

COMMON PREDICTION

New physics at the weak scale


Expectations



Reality



Two Kinds of Physicists



Model-building




EFT

General Idea

Higgs mass:



Production and decay:



MODEL-INDEPENDENT STATEMENT?

Naturalness \longleftrightarrow Higgs couplings

Outline

1. The Ingredients

2. The Recipe

3. The Cake

Outline

1. The Ingredients

2. The Recipe

3. The Cake

The Question

Suppose there is a deviation in hWW or hZZ .
What can we say about new bosons?

Proof by Contradiction

How to bound new bosons?

- ▶ BSM fermions in generic representation
- ▶ Yukawas $\mathcal{L} \supset -yLHN^c - y^c L^c H^\dagger N$



[Arkani-Hamed, Blum, D'Agnolo, Fan 2012; Blum, D'Agnolo, Fan 2015]

Proof by Contradiction

How to bound new bosons?

- ▶ BSM fermions in generic representation
- ▶ Yukawas $\mathcal{L} \supset -yLHN^c - y^c L^c H^\dagger N$
 1. **Deviations** in hWW and hZZ
 2. **Instability** in RGEs
- ▶ Vary masses and representations
- ▶ **Conservative upper bound** on Λ_B



[Arkani-Hamed, Blum, D'Agnolo, Fan 2012; Blum, D'Agnolo, Fan 2015]


Focus on Fermions

Generic requirements:

1. Responsible for **deviations**
2. Consistent with **experiment**
3. No gauge **anomalies**


VECTOR-LIKE FERMIONS

Why These Couplings?



[de Blas et al. 2022]

Why These Couplings?



tree SM vs loop BSM

[de Blas et al. 2022]

Why Beyond EFT?

EFT intuition:

- ▶ Deviation $\delta g_h \simeq c(v^2/\Lambda^2)$
- ▶ New physics $\Lambda \simeq v\sqrt{c/\delta g_h}$
- ▶ **Additionally**, $\Lambda_B > \Lambda$

SPOILER

$\Lambda_B \simeq \Lambda$ for HL-LHC deviations


Outline

1. The Ingredients

2. The Recipe

3. The Cake


Step One



- ▶ Renormalize SM + VLFs $\longrightarrow \delta\mu_{hVV} \sim f(y, M)$
- ▶ Measured quantities α, M_Z, G_F

$$\delta\mu_{hWW} = \frac{y^2 v^2}{80\pi^2 M^2} \left[y^2 \frac{(1 + c_W^2)}{c_W^2 - s_W^2} - \frac{m_h^2}{v^2} - \frac{g^2}{4} \frac{s_W^2}{c_W^2 - s_W^2} \right] + \dots$$

Naive Dimensional Analysis




$$\delta g_{hWW} \sim \frac{g^2 y^4 v^3}{16\pi^2 M^2}$$


Step Two

Two instabilities:

- ▶ Higgs quartic, $d\lambda/d \log \mu \sim -y^4/16\pi^2$
 - ▶ Landau pole, $dy/d \log \mu \sim y^3/16\pi^2$
- } Λ_B
- ▶ **Crucially** only rectified by bosons
 - ▶ SM vacuum decay at $\lambda(\Lambda_B)^{-1} = -14.53 + 0.153 \log(\text{GeV}/\Lambda_B)$




[Coleman 1977; Buttazzo et al. 2013]



Step Two

Two instabilities:

- ▶ Higgs quartic, $d\lambda/d \log \mu \sim -y^4/16\pi^2$
 - ▶ Landau pole, $dy/d \log \mu \sim y^3/16\pi^2$
- } Λ_B
- ▶ **Crucially** only rectified by bosons
 - ▶ SM vacuum decay at $\lambda(\Lambda_B)^{-1} = -14.53 + 0.153 \log(\text{GeV}/\Lambda_B)$



Scaling with Color

$$\delta g_{hVV} \sim r \frac{\alpha_V y^4 v^2}{4\pi M^2} = \frac{\alpha_V y_r^4 v^2}{4\pi r M^2} \lesssim \frac{\alpha_V y_r^4 v^2}{4\pi r M_{\text{exp}}^2}$$


$$\frac{dy}{d \log \mu} \sim r \frac{y^3}{16\pi^2} \longrightarrow y_r \equiv y\sqrt{r} \quad \text{'t Hooft coupling}$$

VECTOR-LIKE LEPTONS

$$L = (1, n)_{Y-1/2}, \quad N^c = (1, n-1)_{-Y}$$

$$L^c = (1, n)_{-Y+1/2}, \quad N = (1, n-1)_Y$$



Step Three



Experimental constraints (EWPTs, collider)

[The Gfitter Group 2018]

Step Three



Experimental constraints (EWPTs, collider)

[The Gfitter Group 2018]

Aside on Strong Coupling

Strong coupling:

$$\text{correction } \mathcal{O}(v^2/f^2), \quad m_* = 4\pi f$$

Compare with

$$\text{correction } \mathcal{O}(y^4 v^2/16\pi^2 M^2) \longrightarrow f_{\text{eff}} \simeq 4\pi M/y^2$$

Outline


1. The Ingredients

2. The Recipe



3. The Cake

Money Plots: hWW


doublet + singlet w/ $Q = 0$, $y = y^c$




Money Plots: hWW



Money Plots: hWW



Money Plots: hZZ



Summary and Future

- ▶ Deviation at HL-LHC **requires** new bosons
- ▶ Deviation at future colliders:
 1. light **new fermions**
 2. **new bosons** under 100 TeV
- ▶ Systematically study $h\gamma\gamma$ and $hZ\gamma$

Summary and Future

- ▶ Deviation at HL-LHC **requires** new bosons
- ▶ Deviation at future colliders:
 1. light **new fermions**
 2. **new bosons** under 100 TeV
- ▶ Systematically study $h\gamma\gamma$ and $hZ\gamma$
- ▶ Powerful probes of new physics!

Thank you!