

Title: General Relativity for Cosmology Lecture - 111623

Speakers: Achim Kempf

Collection: General Relativity for Cosmology

Date: November 16, 2023 - 2:00 PM

URL: <https://pirsa.org/23110004>

Abstract: Zoom: <https://pitp.zoom.us/j/91640855624?pwd=dWVWV2doSnBhUS9JUkhjQVBwY0h0dz09>

GR for Cosmology, Achim Kempf

Lecture 19

Classification of solutions of GR

(and along the way we will introduce some generally useful methods of group theory)

Recall:

- The task is to solve the equations of motion of matter, jointly with the Einstein equation:

$$G_{\mu\nu} = 8\pi G T_{\mu\nu}$$

- In practice, this problem must be simplified, i.e., the number of to-be-determined functions must be reduced.

→ Make symmetry assumptions.

Recall: Symmetries & Killing vector fields

- Two spacetimes $(M, g), (\tilde{M}, \tilde{g})$ are isometric (and therefore of exactly identical shape) if there is a diffeomorphism $\phi: M \rightarrow \tilde{M}$ so that the image of the metric g in \tilde{M} is \tilde{g} : $T_g = \tilde{g}$.
- A space-time has a symmetry, if we find such a ϕ for $\tilde{M} = M$.

Question: How much can we weaken the symmetry assumptions of Friedmann-Lemaitre and still get exact solutions?

Strategy:

- Classify cosmological models $(M, g), T_{\mu\nu}$ by the amount and type of symmetry assumed.
- For each amount and type of symmetry assumed, try to find exact solutions or at least (asymptotic) properties of exact solutions.

Remark: Among the high symmetry models, some come arbitrarily close to F.L. at finite times!

See, e.g., text by Wainwright & Ellis.

Note: The set of all symmetries of a manifold (M, g) forms a "group":

Definition: A "group" G is a set, with an operation, say " \circ ",

$$\circ: G \times G \rightarrow G$$

and a "neutral element", say " e ", $e \in G$, such that

$$(a \circ b) \circ c = a \circ (b \circ c) \quad \forall a, b, c \in G$$

$$a \circ e = e \circ a = a \quad \forall a \in G$$

$$\exists a': a' \circ a = a \circ a' = e \quad \begin{matrix} \uparrow \text{"there exists"} \\ \text{"for all"} \end{matrix}$$

- The task is to solve the equations of motion of matter, jointly with the Einstein equation:

$$G_{\mu\nu} = 8\pi G T_{\mu\nu}$$

- In practice, this problem must be simplified, i.e., the number of to-be-determined functions must be reduced.




→ Make symmetry assumptions.

Recall: Symmetries & Killing vector fields

- Two spacetimes (M, g) , (\tilde{M}, \tilde{g}) are isometric (and therefore of exactly identical shape) if there is a diffeomorphism $\phi: M \rightarrow \tilde{M}$ so that the image of the metric g in \tilde{M} is $\tilde{g}: T_g = \tilde{g}$.

- A space-time has a symmetry, if we find such a ϕ for $\tilde{M} = M$.

- Example:



ϕ performs a rotation of M about a symmetry axis, to obtain $\tilde{M} = M$ with $T_g = \tilde{g}$.

the amount and type of symmetry assumed.

- For each amount and type of symmetry assumed, try to find exact solutions or at least (asymptotic) properties of exact solutions.

Remark: Among the high symmetry models, some come arbitrarily close to F.L. at finite times!

See, e.g., text by Wainwright & Ellis.

Note: The set of all symmetries of a manifold (M, g) forms a "group":

Definition: A "group" G is a set, with an operation, say " \circ ",

$$\circ: G \times G \rightarrow G$$

and a "neutral element", say " e ", $e \in G$, such that

$$(a \circ b) \circ c = a \circ (b \circ c) \quad \forall a, b, c \in G$$

$$a \circ e = e \circ a = a \quad \forall a \in G$$

$$\exists a': a' \circ a = a \circ a' = e \quad \forall a \in G$$

↑ "there exists"

↑ "for all"


Definition: A group G is called a Lie group if G is also a finite-dimensional smooth manifold.

Example: The sets of rotations in \mathbb{R}^3 forms a 3-dimensional Lie group, $SO(3)$.

The angles α, β, γ are coordinates for elements $g \in SO(3)$.

◻ A space-time has a symmetry, if we find such a ϕ for $\tilde{M} = M$.

◻ Example:



ϕ performs a rotation of M about a symmetry axis, to obtain $\tilde{M} = M$ with $T_g = \tilde{g}$.

- Remarks:
- ◻ The symmetries of a manifold (M, g) can be discrete, such as reflections.
 - ◻ But often, the symmetry group of a manifold (M, g) is actually a Lie group.

Note: ◻ Each $h \in G$ yields an isometric diffeomorphism, by assumption.
 $h: M \rightarrow M$, namely $h: p \mapsto h(p) \quad \forall p \in M$

◻ Consider the set $O_p \subset M$ defined by: $O_p = \{q \in M \mid \exists h \in G: h(p) = q\}$

Definition: The set O_p is called the *Orbit* of p under the action of the group G .

Note: If G is a Lie group then each orbit O_p is σ or a submanifold of (M, g) .

Question: What are the infinitesimal isometric diffeomorphisms?
And what type of mathematical structure do the infinitesimal symmetries form?

◻ always for Γ, g compatibility

$$a \circ e = e \circ a = a \quad \forall a \in G$$

$$\exists a': a' \circ a = a \circ a' = e \quad \begin{matrix} \uparrow \text{"there exists"} \\ \text{"for all"} \end{matrix}$$

Definition: A group G is called a *Lie group* if G is also a finite-dimensional smooth manifold.

Example: The sets of rotations in \mathbb{R}^3 forms a 3-dimensional Lie group, $SO(3)$.
The angles α, β, γ are coordinates for elements $g \in SO(3)$.

◻ Recall: The Lie derivative,

$$\begin{aligned} L_{\xi} Q^{a \dots b} &= Q^{a \dots b} \cdot \delta_{j \dots k} \xi^k \\ &= Q^{a \dots b} \cdot \xi_j \delta_{jk} - \dots - Q^{a \dots b} \cdot \xi_k \delta_{jk} \\ &\quad + Q^{a \dots b} \cdot \delta_{j \dots k} \xi^k + \dots + Q^{a \dots b} \cdot \delta_{i \dots k} \xi^k \end{aligned}$$

yields the rate of change of a tensor Q along the flow of diffeomorphisms ϕ generated by a vector field ξ .

⇒ Here, can use L_{ξ} to differentiate along symmetry group orbits.

◻ Thus, if $L_{\xi} g_{\mu\nu} = 0$

then ξ generates isometries $\phi: M \rightarrow M, g \mapsto \tilde{g} = g$.

Note: □ Each $h \in G$ yields an isometric diffeomorphism, by assumption.

$$h: M \rightarrow M, \text{ namely } h: p \mapsto h(p) \quad \forall p \in M$$

□ Consider the set $O_p \subset M$ defined by: $O_p = \{q \in M \mid \exists h \in G: h(p) = q\}$

Definition: The set O_p is called the **Orbit** of p under the action of the group G .

Note: If G is a Lie group then each orbit O_p is par a submanifold of (M, g) .

Question: What are the infinitesimal isometric diffeomorphisms?

And what type of mathematical structure do the infinitesimal symmetries form?

$$\square \text{ But } L_g g_{\mu\nu} = \overset{\text{if always for } \Gamma, g \text{ compatible}}{\cancel{g^k}} g_{\mu\nu;k} + g_{\mu k} \overset{\text{if }}{\cancel{g^k}}{}_{;k} + g_{\nu k} \overset{\text{if }}{\cancel{g^k}}{}_{;k}$$

\Rightarrow A vector field ξ generates a symmetry of spacetime if it is a **Killing vector field**:

$$\xi_{\mu;\nu} + \xi_{\nu;\mu} = 0 \quad (\times)$$

Q: Maximum number, d , of Killing vector fields in n dims?

A: $d = n(n+1)/2$ To see this, note that there are 2 ways to obey Eq. (1):

a) $\xi_{\mu;\nu} = 0 \quad \forall \nu$, i.e. $\nabla \xi = 0$

(can have maximally n such indep. vectors)

b) $\nabla \xi \neq 0$, but then $K_{\mu\nu} := \xi_{\mu;\nu}$ is antisymmetric

(can have at most $n(n-1)/2$ indep. such cases.)

$$\left. \begin{aligned} d &= n + \\ &n(n-1)/2 \\ &n(n+1)/2 \end{aligned} \right\}$$

$$+ Q^{(n)}_{\mu_1 \dots \mu_n} \xi^{\mu_1} + \dots + Q^{(n)}_{\mu_1 \dots \mu_n} \xi^{\mu_n}$$

yields the rate of change of a tensor Q along the flow of diffeomorphisms ϕ generated by a vector field ξ .

\Rightarrow Here, can use L_ξ to differentiate along symmetry group orbits.

□ Thus, if $L_\xi g_{\mu\nu} = 0$

then ξ generates isometries $\phi: M \rightarrow M, g \mapsto \tilde{g} = g$.

From a symmetry Lie group to a "symmetry Lie algebra":

General idea:

Normally the points of a manifold cannot be multiplied!

- A Lie group is a smooth manifold with extra structure: the multiplication.
- Notion: Product of group elements close to **1** yields a group element close to **1**.
- Consider the tangent space $T_e(G)$ to the point **1** of the Lie group manifold **G**.
- $T_e(G)$ is a vector space and it has extra structure, inherited from the group's multiplication.
- Define the **Lie algebra** of a group G to be $T_e(G)$, equipped with the inherited "multiplication".

The identity element of the group, $e = 1$ is also a point of the group's manifold. $T_e(G)$ is the tangent space to this point.

Crucial fact: From knowledge of only the Lie algebra, i.e., only $T_e(G)$ and its "multiplication", the group **G** can be constructed!

(though not always unique)

Q: Maximum number, d , of Killing vector fields in n dims?

A: $d = n(n+1)/2$ To see this, note that there are 2 ways to obey Eq. (1):

a) $\xi_{\mu;\nu} = 0 \quad \forall \nu$, i.e. $\nabla \xi = 0$
 can have maximally n such indep. vectors

b) $\nabla \xi \neq 0$, but then $K_{\mu\nu} := \xi_{\mu;\nu}$ is antisymmetric
 can have at most $n(n-1)/2$ indep. such cases.

$$\begin{aligned} d &= n + \\ &n(n-1)/2 \\ &n(n+1)/2 \end{aligned}$$

Let us collect the properties that the inherited multiplications of all Lie algebras share.

Then, let us define **Lie algebra** as anything with these properties:

Definition:

A Lie algebra is a vector space A , with an operation $\{\cdot, \cdot\}$

$$\{\cdot, \cdot\}: A \times A \rightarrow A \quad \text{"Lie bracket"}$$

obeying $\{\tau, s\} = -\{s, \tau\} \quad \forall \tau, s \in A$
 "Jacobi identity"
 and $\{\{\tau, s\}, t\} + \{\tau, \{\tau, s\}\} + \{\{\tau, s\}, t\} = 0$

Theorem: Every vector space A with a "multiplication" $\{\cdot, \cdot\}$ that obeys these axioms is isomorphic to $T_p(G)$ of a Lie group G .

Consider the tangent space $T_p(G)$ to the point $p \in G$ of the Lie group manifold G .

$T_p(G)$ is a vector space and it has extra structure, inherited from the group's multiplication.

Define the Lie algebra of a group G to be $T_p(G)$, equipped with the inherited "multiplication".

The identity element of the group, $p = 1$ is also a point of the group's manifold.
 $T_p(G)$ is the tangent space to this point.

Crucial fact: From knowledge of only the Lie algebra, i.e., only $T_p(G)$ and its "multiplication", the group G can be constructed!
 (though not always uniquely)

Proposition: The set of Killing vector fields $\xi^{(i)}$ of (M, g) is a Lie algebra.

Exercise: Prove this, i.e., show the following:

Assume $\xi^{(1)}, \xi^{(2)}$ are Killing vector fields of (M, g) and $\alpha, \beta \in \mathbb{R}$.

Then: $\alpha \xi^{(1)} + \beta \xi^{(2)}$ (i.e., they form a vector space)

$$\text{and } \{\xi^{(1)}, \xi^{(2)}\} := \xi^{(1)} \xi^{(2)} - \xi^{(2)} \xi^{(1)}$$

are also Killing vector fields,

and the $\xi^{(i)}$ obey the Jacobi identity.

\Rightarrow A vector field ξ generates a symmetry of spacetime if it is a Killing vector field:

$$\xi_{\mu;\nu} + \xi_{\nu;\mu} = 0 \quad (\text{X})$$

Q: Maximum number, d , of Killing vector fields in n dims?

A: $d = n(n+1)/2$ To see this, note that there are 2 ways to obey Eq. (X):

a) $\xi_{\mu;\nu} = 0 \forall \nu$, i.e. $\nabla \xi = 0$
 can have maximally n such indep. vectors

b) $\nabla \xi \neq 0$, but then $K_{\mu\nu} := \xi_{\mu;\nu}$ is antisymmetric
 can have at most $n(n-1)/2$ indep. such cases.

$$\begin{aligned} d &= n + \\ &n(n-1)/2 \\ &= n(n+1)/2 \end{aligned}$$

- Let us collect the properties that the inherited multiplications of all Lie algebras share.
- Then, let us define Lie algebras as anything with these properties:

Definition:

A Lie algebra is a vector space A , with an operation $\{\cdot, \cdot\}$

$$\{\cdot, \cdot\}: A \times A \rightarrow A \quad \text{"Lie bracket"}$$

obeying $\{\tau, s\} = -\{s, \tau\} \quad \forall \tau, s \in A$
 "Jacobi identity"
 and $\{\{\tau, s\}, t\} + \{\tau, \{s, t\}\} + \{s, \{\tau, t\}\} = 0$

General idea:

Normally the points of a manifold cannot be multiplied!

- A Lie group is a smooth manifold with extra structure: the multiplication.
- Notion: Product of group elements close to 1_G yields a group element close to 1 .
- Consider the tangent space $T_1(G)$ to the point 1_G of the Lie group manifold G .
- $T_1(G)$ is a vector space and it has extra structure, inherited from the group's multiplication.
- Define the Lie algebra of a group G to be $T_1(G)$, equipped with the inherited "multiplication".

The identity element of the group, $p = 1$, is also a point of the group's manifold. $T_1(G)$ is the tangent space to this point.

Crucial fact: From knowledge of only the Lie algebra, i.e., only $T_1(G)$ and its "multiplication", the group G can be constructed!
 (though not always uniquely)

Proposition: The set of Killing vector fields $\xi^{(1)}$ of (M, g) is a Lie algebra.

Exercise: Prove this, i.e., show the following:

Assume $\xi^{(1)}, \xi^{(2)}$ are Killing vector fields of (M, g) and $\alpha, \beta \in \mathbb{R}$.

Then: $\alpha \xi^{(1)} + \beta \xi^{(2)}$ (i.e., they form a vector space)

$$\text{and } \{\xi^{(1)}, \xi^{(2)}\} := \xi^{(1)} \xi^{(2)} - \xi^{(2)} \xi^{(1)}$$

$$\xi_{\mu;\nu} + \xi_{\nu;\mu} = 0 \quad (\text{X})$$

Q: Maximum number, d , of Killing vector fields in n dims?

A: $d = n(n+1)/2$ To see this, note that there are 2 ways to obey Eq. (X):

a) $\xi_{\mu;\nu} = 0 \quad \forall \nu$, i.e. $\nabla \xi = 0$

(can have maximally n such indep. vectors)

b) $\nabla \xi \neq 0$, but then $K_{\mu\nu} := \xi_{\mu;\nu}$ is antisymmetric

(can have at most $n(n-1)/2$ indep. such cases.)

$$\left. \begin{array}{l} d = n + \\ n(n-1)/2 \\ n(n+1)/2 \end{array} \right\} \Rightarrow$$

□ Let us collect the properties that the inherited multiplications of all Lie algebras share.

□ Then, let us define **Lie algebras** as anything with these properties:

Definition:

A Lie algebra is a vector space A , with an operation $\{\cdot, \cdot\}$

$$\{\cdot, \cdot\}: A \times A \rightarrow A \quad \text{"Lie bracket"}$$

obeying $\{\tau, s\} = -\{s, \tau\} \quad \forall \tau, s \in A$
 "Jacobi identity"

and $\{\{\tau, s\}, t\} + \{\{\tau, r\}, s\} + \{\{s, r\}, \tau\} = 0$

Theorem: Every vector space A with a "multiplication" $\{\cdot, \cdot\}$ that obeys these axioms is isomorphic to $T_x(G)$ of a Lie group G .

□ Notice: Product of group elements close to 1_G yields a group element close to 1 .

□ Consider the tangent space $T_x(G)$ to the point 1_G of the Lie group manifold G .

□ $T_x(G)$ is a vector space and it has extra structure, inherited from the group's multiplication.

□ Define the Lie algebra of a group G to be $T_x(G)$, equipped with the inherited "multiplication".

The identity element of the group, $p=1$, is also a point of the group's manifold. $T_x(G)$ is the tangent space to this point.

Crucial fact: From knowledge of only the Lie algebra, i.e., only $T_x(G)$ and its "multiplication", the group G can be constructed! (though not always uniquely)

Proposition: The set of Killing vector fields $\xi^{(i)}$ of (M, g) is a Lie algebra.

Exercise: Prove this, i.e., show the following:

Assume $\xi^{(1)}, \xi^{(2)}$ are Killing vector fields of (M, g) and $\alpha, \beta \in \mathbb{R}$.

Then: $\alpha \xi^{(1)} + \beta \xi^{(2)}$ (i.e., they form a vector space)

$$\text{and } \{\xi^{(1)}, \xi^{(2)}\} := \xi^{(1)}\xi^{(2)} - \xi^{(2)}\xi^{(1)}$$

are also Killing vector fields,

and the $\xi^{(i)}$ obey the Jacobi identity.

$\{, \}$: $A \times A \rightarrow A$ "Lie bracket"
 obeying $\{r, s\} = -\{s, r\} \quad \forall r, s \in A$
 and $\{\{r, s\}, t\} + \{s, \{r, t\}\} + \{r, \{s, t\}\} = 0$

Theorem: Every vector space A with a "multiplication" $\{, \}$ that obeys these axioms is isomorphic to $T_e(G)$ of a Lie group G .

Summary of the big picture:

1. The symmetries of any (M, g) form a group: they can be concatenated associatively, and all possess an inverse. Some symmetries are differentiable, parametrized by the flow \Rightarrow the symmetries form a Lie group.
2. Each Killing vector field is the infinitesimal generator of a flow of isometric diffeomorphisms, i.e., of a symmetry.
3. We see here that the Killing vector fields indeed form a Lie algebra.
4. Recall that every Lie algebra generates a Lie group.

Assume ξ, ζ are Killing vector fields of (M, g) and $\alpha, \beta \in \mathbb{R}$.

Then: $\alpha \xi^{(1)} + \beta \xi^{(2)}$ (i.e., they form a vector space)
 and $\{\xi^{(1)}, \xi^{(2)}\} := \xi^{(1)} \xi^{(2)} - \xi^{(2)} \xi^{(1)}$
 are also Killing vector fields,
 and the $\xi^{(i)}$ obey the Jacobi identity.

Surfaces of homogeneity and the isotropy subgroup:

□ Definition:

Let r be the dimension of the Lie algebra, i.e., also the dimension of the Lie group of symmetries.

□ Recall this definition:

- Consider the set of points $\Omega(p)$ that a point p can flow to along the Killing vector fields.
- $\Omega(p)$ is called the orbit of $p \in M$ under the action of the symmetry group. We denote the dimension of the orbit by s .

form a Lie group.

2. Each Killing vector field is the infinitesimal generator of a flow of isometric diffeomorphisms, i.e., of a symmetry.
3. We see here that the Killing vector fields indeed form a Lie algebra.
4. Recall that every Lie algebra generates a Lie group.

□ Clearly:

The dimension of an orbit cannot be larger than the dimension of the symmetry group, i.e.

$$s \leq r,$$

but $s < r$ easily happens:

□ Example:

- Consider $M := \mathbb{R}^2$ and $p = (0,0)$.
- Then $r = r_{\max} = \sqrt{n(n+1)/2} = \underline{\underline{3}}$ is dim. of sym. group.
- \Rightarrow The three-dimensional Lie algebra of Killing vector fields is spanned by three Killing vector fields:

□ Recall this definition:

- Consider the set of points $O(p)$ that a point p can flow to along the Killing vector fields.
- $O(p)$ is called the orbit of $p \in M$ under the action of the symmetry group. We denote the dimension of the orbit by s .

□ Concretely:

$$K^{(1)} := \frac{\partial}{\partial x}, \quad K^{(2)} := \frac{\partial}{\partial y}$$

$$K^{(3)} := Y \frac{\partial}{\partial x} - X \frac{\partial}{\partial y}$$

Group elements generated by them
 $e^{\frac{\partial}{\partial x} + b \frac{\partial}{\partial y}}$ and they act as
 $e^{\frac{\partial}{\partial x} + b \frac{\partial}{\partial y}} f(x,y) = f(x+a, y+b)$
by Taylor expansion.

□ Orbit of $p = (0,0)$:

$$O(p) = \mathbb{R}^2 \text{ because generators } \frac{\partial}{\partial x} \text{ and } \frac{\partial}{\partial y} \text{ generate flow to every where.}$$

Def: The surface of homogeneity has dimension $s=2 < r$
generated by the Killing vectors (here: $K^{(1)}, K^{(2)}$) which do not have closed orbits

- Notice: Since $n=2$, at any given point p , only at most 2 Killing vectors can be linearly independent at p .

but $s < r$ easily happens:

Example:

- 1 Consider $M = \mathbb{R}^2$ and $p = (0,0)$.
- 2 Then $r = r_{\max} = \sqrt{n(n+1)/2} = \underline{\underline{3}}$ is dim. of sym. group.
- 3 \Rightarrow The three-dimensional Lie algebra of Killing vector fields is spanned by three Killing vector fields:

Rôle of $K^{(3)}$?

$(K^{(3)})$ is the angular momentum
and it of course generates rotations:
 $e^{-itK^{(3)}}f(x,y) = f(x\cos t - y\sin t, x\sin t + y\cos t)$

The flow generated by $K^{(3)}$ leaves p fixed and rotates everything around p .

Definition:

We say that those Killing vector fields which do not generate a homogeneity surface, i.e., which generate a trivial group orbit for a point are generating the isotropy subgroup (of the full symmetry group generated by all Killing vectors).

Dimension, d , of the isotropy subgroup?

Clearly: $d = r - s$

isotropic full homogeneous

Orbit of $p = (0,0)$:

$O(p) = \mathbb{R}^2$ because generators $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ generate flow to everywhere.

$e^{t\frac{\partial}{\partial x}}f(x,y) = f(x+t, y)$
by Taylor expansion.

Def: The surface of homogeneity has dimension $s = 2 < r$

generated by the Killing vectors (here: $K^{(3)}, K^{(1)}$) which do not have fixed orbits

Notice: Since $n=2$, at any given point p , only at most 2 Killing vectors can be linearly independent at p .

Classification of cosmological models

The classification is with respect to:

Dimension of isotropy subgroup d :

(# of conserved angular momenta) \rightarrow
 $d = 0, 1, 2, 3, 4, 5, 6$
 e.g. full Lorentz symmetry
 anisotropic case
 at each Poincaré one rotational symmetry axis
 e.g. spatially isotropic case

Dimension of homogeneity surfaces s :

(# of conserved momenta) \rightarrow
 $s = 0, 1, 2, 3, 4$
 inhomogeneous
 homogeneous
 homogeneous on 3-dimensional

which do not generate a homogeneity surface,
 i.e., which generate a trivial group orbit for a point
 are generating the isotropy subgroup (of the
 full symmetry group generated by all Killing vectors).

Dimension, d , of the isotropy subgroup?

Clearly: $d = r - s$



isotropy full homogeneous.

A large body of literature exists on most cases of (d, s) :

- Many exact solutions are known!
- Many asymptotic behaviors are known!
- Comprehensive text:

Wainwright & Ellis, Dyn. systems in cosmology,
 Cambridge Univ. Press (1997)

Examples:


α	homogeneity ↓	isotropy ↓
4	5	d
4	3	Einstein's static model
4	1	Gödel's model
4	0	Ozsváth-Kovács models
3	3	Friedmann-Lemaitre models
3	1	spatially hom & locally one rot. symm axis
3	0	Bianchi models
:	:	

Definition: u is called the "fundamental 4-velocity field"

anisotropic case
 at each point
 one rotational
 symmetry axis

e.g. spatially
 isotropic case

Dimension of homogeneity surfaces s :



Powerful alternative classification approach:

Idea: Classify the possible $T_{\mu\nu}$, then use Einstein equation to obtain classification of curvature.

Proposition:

For every physical energy momentum tensor $T_{\mu\nu}$ there exists a unique timelike vector field u so that $T_{\mu\nu}$ takes this standard form:

$$T_{ab} = \overset{\text{scalar}}{\mu u_a u_b} + \overset{\text{vector}}{q_a u_b + q_b u_a} + \overset{\text{scalar}}{\rho(g_{ab} + u_a u_b)} + \overset{\text{tensor}}{\pi_{ab}}$$

where q and π are a vector field and a tensor field obeying:

$$q_a u^a = 0, \quad \pi_{ab} u^b = 0, \quad \pi_a^a = 0, \quad \pi_{ab} = \pi_{ba}$$

Sasaki classification:

II Comprehensive text:

Wainwright & Ellis, Dyn. systems in cosmology,
Cambridge Univ. Press (1997)

Sharing Desktop ▾ Stop Sharing

Examples:

homogeneity ↓ isotropy ↑

a) s d


- 4 3 Einstein's static model
- 4 1 Gödel's model 
- 4 0 Ozerath-Kerr models
- 3 3 Friedmann-Lemaître models
- 3 1 spatially homogeneous & locally one rot. sym. axis
- 3 0 Bianchi models
- :
- :

Definition: u is called the "fundamental 4-velocity field"Note: E.g., for a perfect fluid this is the fluid velocity:

$$T_{ab} = \mu u_a u_b + p(g_{ab} + u_a u_b), \quad u_a u^a = -1$$

Recall: equation of state is

$$p = (\gamma - 1)\mu$$

$$\gamma = \begin{cases} 1 & \text{dust} \\ 4/3 & \text{radiation} \\ 0 & \text{cosmological constant} \end{cases}$$


II Definition:

If (M, g) possesses spacelike $S=3$ homogeneity but the fundamental velocity is not orthogonal to the homogeneity surfaces, then we say that this cosmology is "tilted".

Proposition:

For every physical energy momentum tensor $T_{\mu\nu}$ there exists a unique timelike vector field u so that $T_{\mu\nu}$ takes this standard form:

$$T_{ab} = \frac{\text{scalar}}{\text{scalar}} u^a u_b + \frac{\text{vector}}{\text{vector}} u_a u^b + \frac{\text{scalar}}{\text{scalar}} g_{ab} + p(g_{ab} + u_a u_b) + \pi_{ab}$$

where q and π are a vector field and a tensor field obeying:

$$q_a u^a = 0, \quad \pi_{ab} u^b = 0, \quad \pi_a^a = 0, \quad \pi_{ab} = \pi_{ba}$$

Segré classification:

I A systematic classification of $T_{\mu\nu}$ can be performed, by the analysis of its eigenvalues/eigenvectors. Nontrivial because:

II $T_{\mu\nu}$ is symmetric.But, the inner product in the vector space is $g_{\mu\nu} \Rightarrow T_{\mu\nu}$ is generally not hermitian!

III $T_{\mu\nu}$ is in a space with the inner product $g^{\mu\nu} = \delta^{\mu\nu}$, but $T^{\mu\nu}$ is generally not symmetric!


Use Jordan normal form:

⇒ Segre classification yields 4 main types of energy momentum tensors $T_{\mu\nu}$

Recall: equation of state is

$$p = \tilde{\mu}(\tilde{e}-1)\mu$$

$$\tilde{\mu} = \begin{cases} 1 & \text{dust} \\ 4/3 & \text{radiation} \\ 0 & \text{cosmological constant} \end{cases}$$



□ Definition:

If (M, g) possesses spacelike $s=3$ homogeneity but the fundamental velocity is not orthogonal to the homogeneity surfaces, then we say that this cosmology is "tilted".

Recall strategy:

The classification of possible $T_{\mu\nu}$ should, via the Einstein eqns, yield a classification of possible curvatures.

Indeed: In 3+1 dimensions the Einstein equation also reads:

$$R_{\mu\nu} = 8\pi G (T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T)$$

Exercise: Prove this and notice the dimension-dependence

\Rightarrow The 10 degrees of freedom of $T_{\mu\nu}$ (as a symmetric 4x4 matrix) determine the 10 degrees of freedom of $R_{\mu\nu}$.

But, the inner product in the vector space is $g_{\mu\nu}$ $\Rightarrow T_{\mu\nu}$ is generally not hermitian!

□ T^*_{ν} is in a space with the inner product $g^{*\nu} = g^{\nu}$, but T^*_{ν} is generally not symmetric!

Use Jordan normal form:

\Rightarrow Segré classification yields 4 main types of energy momentum tensors $T_{\mu\nu}$.

\Rightarrow The Segré classification of possible $T_{\mu\nu}$ yields, via the Einstein equation also a classification of possible Ricci tensors $R_{\mu\nu}$.

Q: Does this yield also a classification of the possible Riemann tensors $R^*_{\nu\mu\rho}$?

A: No! The Ricci tensor contains only 10 of the 20 degrees of freedom of the Riemann tensor! (In 3+1 dim)

Prop.: The information in $R^*_{\nu\mu\rho}$ is shared among the Ricci tensor $R_{\mu\nu}$ and the so-called Weyl tensor, $C^*_{\nu\mu\rho}$.

\Rightarrow It remains to classify the possible Weyl tensors!

Indeed: In 3+1 dimensions the Einstein equation also reads:

$$R_{\mu\nu} = 8\pi G (T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T)$$

Exercise: Prove this and notice the dimension-dependence

\Rightarrow The 10 degrees of freedom of $T_{\mu\nu}$ (as a symmetric 4x4 matrix) determine the 10 degrees of freedom of $R_{\mu\nu}$.

The Weyl tensor, $C^{\alpha\beta}_{\gamma\eta}$:

$$C^{\alpha\beta}_{\gamma\eta} := R^{\alpha\beta}_{\gamma\eta} - \frac{1}{2}(g^{\alpha}_{\gamma}R^{\beta}_{\eta} + g^{\beta}_{\eta}R^{\alpha}_{\gamma} - g^{\alpha}_{\eta}R^{\beta}_{\gamma} - g^{\beta}_{\gamma}R^{\alpha}_{\eta}) + \frac{1}{6}(g^{\alpha}_{\gamma}g^{\beta}_{\eta} - g^{\alpha}_{\eta}g^{\beta}_{\gamma})R$$

Notice: If R^{α}_{β} and $C^{\alpha\beta}_{\gamma\eta}$ are given, they determine $R^{\alpha\beta}_{\gamma\eta}$ fully:

$$\begin{aligned} R^{\alpha\beta}_{\gamma\eta} &= C^{\alpha\beta}_{\gamma\eta} + \frac{1}{2}(g^{\alpha}_{\gamma}R^{\beta}_{\eta} + g^{\beta}_{\eta}R^{\alpha}_{\gamma} \\ &\quad - g^{\alpha}_{\eta}R^{\beta}_{\gamma} - g^{\beta}_{\gamma}R^{\alpha}_{\eta}) - \frac{1}{6}(g^{\alpha}_{\gamma}g^{\beta}_{\eta} - g^{\alpha}_{\eta}g^{\beta}_{\gamma})R \end{aligned}$$

$\Rightarrow R^{\alpha\beta}_{\gamma\eta}$ is expressed through $C^{\alpha\beta}_{\gamma\eta}$ and $R^{\alpha\beta}$

\Rightarrow The Weyl tensor $C^{\alpha\beta}_{\gamma\eta}$ indeed contains all that information about the curvature $R^{\alpha\beta}_{\gamma\eta}$, which is not in $R^{\alpha\beta}_{\gamma\eta}$.
Determined from $T_{\mu\nu}$ via the Einstein eqn.

possible Riemann tensors $R^{\alpha\beta}_{\gamma\eta}$?

A: No! The Ricci tensor contains only 10 of the 20 degrees of freedom of the Riemann tensor! (In 3+1 dim)

Prop.: The information in $R^{\alpha\beta}_{\gamma\eta}$ is shared among the Ricci tensor $R_{\mu\nu}$ and the so-called Weyl tensor, $C^{\alpha\beta}_{\gamma\eta}$.

\Rightarrow It remains to classify the possible Weyl tensors!

$\Rightarrow C^{\alpha\beta}_{\gamma\eta}$ contains all that curvature information which is not determined via the Einstein equation by $T_{\mu\nu}$.

$\Rightarrow C^{\alpha\beta}_{\gamma\eta}$ describes all that curvature which can exist even where there is no matter! (e.g. gravity waves)

also e.g. sun's gravity away from the sun in empty space

Proposition

Assume (M, g) is a 3+1 dimensional Lorentzian manifold.

Choose any smooth positive scalar function ϕ on M .

Define (M, \tilde{g}) with the new metric \tilde{g} obtained through the "conformal transformation":

$$g_{\mu\nu}(x) \rightarrow \tilde{g}_{\mu\nu}(x) = \phi(x)g_{\mu\nu}(x)$$

Then: $\tilde{C}^{\alpha\beta\gamma\eta}(x) = C^{\alpha\beta\gamma\eta}(x) \quad \forall x \in M$ (Exercise: wh 21/29 would be prof st. 21/29)

$$R^{\mu\nu}_{rs} = C^{\mu\nu}_{rs} + \frac{1}{2} (g^{\mu}_r R^{\nu}_s + g^{\nu}_r R^{\mu}_s - g^{\mu}_s R^{\nu}_r - g^{\nu}_s R^{\mu}_r) - \frac{1}{6} (g^{\mu}_r g^{\nu}_s - g^{\nu}_r g^{\mu}_s) R$$

$\rightsquigarrow R^{\mu\nu}_{rs}$ is expressed through $C^{\mu\nu}_{rs}$ and $R^{\mu\nu}$

↑ 20 indep. components ↑ 10 indep. comp. ↑ 10 indep. comp.

⇒ The Weyl tensor $C^{\mu\nu}_{rs}$ indeed contains all that information about the curvature $R^{\mu\nu}_{rs}$, which is not in $R^{\mu\nu}$.

Determined from $T_{\mu\nu}$ via the Einstein eqn.

Historical remark

- Consider the equivalence class of spacetimes (M, \tilde{g}) that are conformally equivalent to Minkowski space:
- $g_{\mu\nu}(x) = \phi^2(x) \eta_{\mu\nu}$
- Einstein and Fokker initially considered a theory in which the metric possesses only this conformal degree of freedom ϕ (to play role of Newton's gravitational potential).

Newton gravity does come out correctly as a limiting case!

- Then, $S = \int_M R \sqrt{-g} d^4x + \int_M L_{matter} V_f d^4x$ and $\frac{\delta S}{\delta \phi} = 0$ yield: $R = 8\pi G T^M_{\mu\nu}$
- Equivalence principle ok. $T^{(EM)\mu\nu} = 0$
i.e. EM fields would not gravitate.
- Light bending & Mercury perihelion shift wrong.

No gravity wave because
 $C^{\mu\nu}_{\mu\nu} = C^{\mu\nu}_{\nu\nu}$ (Minkowski)
 $= 0$

Proposition

also e.g. sun's gravity away from the sun in empty space

- Assume (M, g) is a 3+1 dimensional Lorentzian manifold.
- Choose any smooth positive scalar function ϕ on M .
- Define (M, \tilde{g}) with the new metric \tilde{g} obtained through the "conformal transformation":

$$g_{\mu\nu}(x) \rightarrow \tilde{g}_{\mu\nu}(x) := \phi(x) g_{\mu\nu}(x)$$

Then: $\tilde{C}^{\mu\nu\rho\sigma}(x) = C^{\mu\nu\rho\sigma}(x) \quad \forall x \in M$ (Exercise: what would happen if $\phi = 0$?)

Recall: via the Einstein equation the Segre classification implies a classification of properties of the Ricci tensor $R_{\mu\nu}$.

It remains to classify the Weyl tensor:

Petrov classification:

This is a classification of the Weyl tensor $C^{\mu\nu}_{\rho\sigma}$, which possesses the 10 remaining degrees of freedom of $R^{\mu\nu}_{\rho\sigma}$.

- $C^{\mu\nu}_{\rho\sigma}$, just like the Riemann tensor, is antisymmetric in $\mu \leftrightarrow \nu$ and in $\rho \leftrightarrow \sigma$, and symmetric in $\mu \leftrightarrow \rho$ & $\nu \leftrightarrow \sigma$.

◻ Einstein and Fokker initially consider theory in which the metric possesses only this conformal degree of freedom (to play role of Newton's gravitational potential).

Newton gravity does come out correctly as a limiting case!

◻ Then, $S = \int_M g_{ij} d^4x + \int_M \text{matter} T_{ij} dx$ and $\frac{\delta S}{\delta g} = 0$ yield:

$$R = 8\pi G T_{\mu\nu}$$

in electromagnetism $T^{(EM)}_{\mu\nu} = 0$
(i.e. EM field would not gravitate.)

No gravity now
but because
 $C_{cd}^{ab} = C_{cb}^{ad}$ (unitarity)
 $= 0$

◻ Equivalence principle ok.

◻ Light bending & Mercury perihelion shift wrong.

◻ Thus $C^{ab}_{\mu\nu}$ can locally be viewed as a symmetric map from the antisymmetric part $A_p(\mathcal{M})^2$ of $T_p(\mathcal{M})^2$ (so called bi-vectors) into itself:

$$G: A_p(\mathcal{M})^2 \rightarrow A_p(\mathcal{M})^2$$

◻ But, the inner product in $A_p(\mathcal{M})^2$ is not positive definite!

$\Rightarrow G$ is generally not hermitian.

Therefore, use Jordan normal form again:

Result: 6 main Petrov classes for Weyl curvature:
according to eigenvalues/eigenvector decomposition.

Type 0: Weyl curvature vanishes

This is a classification of the Weyl tensor $C^{ab}_{\mu\nu}$, which possesses the 10 remaining degrees of freedom of $R^a_{\mu\nu b}$.

◻ $C^{ab}_{\mu\nu}$, just like the Riemann tensor, is antisymmetric in $\mu\nu$ and in ab , and symmetric in $\mu\nu ab$.

Type D: "Static" Weyl curvature, e.g. in vicinity of a star.

Type N: Transverse gravitational waves, the type LIGO detects. Like light, their strength decays $\sim \frac{1}{r}$ from the source.

Type I: Longitudinal gravitational waves.

These waves cause a shear effect.

However, they decay fast: $\sim \frac{1}{r^2}$

Why? Gravitational waves, when small enough, travel with speed of light. Like light, they then cannot oscillate longitudinally.

Types II, III: Mixtures of the above.

But, the inner product in $A_p(\mathbb{M})^*$ is not positive definite!

$\Rightarrow C$ is generally not hermitian.

Therefore, use Jordan normal form again:

Result: 6 main Petrov classes for Weyl curvature:
according to eigenvalues/eigenvalue decomposition.

Type 0: Weyl curvature vanishes

□ Potential problem: (with symmetry assumptions):

(E.g.: recall that flatness
in FL spacetimes is unstable)

□ The so-obtained highly symmetric solutions,
e.g. Friedmann-Lemaître, may possess properties
that are peculiar to high symmetry.

(E.g.:
In Newtonian gravity, a slightly
non-symmetric collapse of a star
would not lead to a singularity
but to a bounce - think fire-starter.)

□ E.g.: When a Friedmann-Lemaître solution,
or a Schwarzschild solution exhibits a
singularity: Is it due to symmetry, or realistic?

□ Singularity theorems (see later) confirm the
robustness under certain conditions
(such as strong energy condition).

→ More confidence in significance of the properties
of highly symmetric solutions.

Type I: Longitudinal gravitational waves.

These waves cause a shear effect.

However, they decay fast: $\sim \frac{1}{r^2}$

Why? Gravitational waves, when small enough, travel with speed of light. Like light, they then cannot oscillate longitudinally.

Types II, III: Mixtures of the above.