Title: Quantum Foundations Lecture

Speakers: Lucien Hardy

Collection: Quantum Foundations

Date: January 24, 2024 - 10:15 AM

URL: https://pirsa.org/24010066

Some questions about any given interpretation.

(0) What is the ontic state?

(1) How does the interpretation solve the measurement problem?

(2) Is the interpetation nonlocal?

(3) Is the interpretation a good departure point for making progress in physics (e.g. solving the problem of Quantum Gravity)?

The de Broglie Bohm model.

(1927) (1952)

Take non-relativistic QM with

$\psi(\vec{x}_1, \vec{x}_2, \cdots \vec{x}_N, t)$ evolves according to

$$i\hbar \frac{\partial \psi}{\partial t} = \sum_{n=1}^{N} -\frac{\hbar^2}{2m_n} \nabla^2 \psi + V\psi$$

$\vec{x}_i$: n-vector in 3 space.

(0) The ontic state at time t is

$$\left(\psi(\vec{x}_1, \vec{x}_2, \ldots \vec{x}_N, t), (\vec{X}_1, \vec{X}_2, \ldots \vec{X}_N)\right)$$

ψ wave fn.

actual positions of actual particles.

$$x = (\vec{x}_1, \vec{x}_2, \ldots \vec{x}_N), \quad X = (\vec{X}_1, \vec{X}_2, \ldots \vec{X}_N)$$

The dynamics is given by the coupled eqns:

(1) $i\hbar \frac{\partial \psi}{\partial t} = \sum_{n=1}^{N} -\frac{\hbar^2}{m_n} \nabla_n^2 \psi + V\psi$ Schrö eqn

(2) $\frac{d\vec{X}_n}{dt} = \frac{\hbar}{m_n} \frac{\mathrm{Im}(\psi^* \nabla_n \psi)}{\psi^* \psi}\Big|_{x=X}$ guidance eqn.

Sometimes an additional postulate is added.

(3) At time $t=0$

$$\rho(\vec{x}_1, \vec{x}_2, \cdots \vec{x}_N, t=0) = |\psi(\vec{x}_1, \vec{x}_2, \cdots \vec{x}_N, t=0)|^2$$

Valentini



What happens when we evolve?

If $\rho = |\psi|^2$ at $t=0$

then eqns (1) and (2) implie that

$\rho = |\psi|^2$ for all later times.

This property is called equivariance.

$\rho = |\psi|^2$ is called equilibrium.

Sometimes an additional postulate is added.

(3) At time $t=0$

$$\rho(\vec{x}_1, \vec{x}_2, \cdots \vec{x}_N, t=0) = |\psi(\vec{x}_1, \vec{x}_2, \cdots \vec{x}_N, t=0)|^2$$

Valentini

$$\vec{J}_n^{(\psi)} = \frac{\hbar}{m_n} \mathrm{Im}(\psi^* \nabla_n \psi)$$

Sketching dBB trajectories.

(1) The no crossing principle.

$$\dot{X} = \left(\dot{\vec{X}}_1, \dot{\vec{X}}_2, \cdots \dot{\vec{X}}_N\right) = f^{\psi}(X, t)$$

no crossing in configuration space.


(X, t)
X_2
not allowed.
X_1

(2) $\rho = |\psi|^2$ at all times.