

Title: Quantum Field Theory for Cosmology - Lecture 20240213

Speakers: Achim Kempf

Collection: Quantum Field Theory for Cosmology (PHYS785/AMATH872)


Date: February 13, 2024 - 4:00 PM

URL: <https://pirsa.org/24020011>

QFT for Cosmology

Project: Simulate quantum field fluctuations

- Task:
- For various important states of scalar QFT, calculate the probability distribution for finding certain outcomes when 'measuring' all $\hat{\phi}(x)$ simultaneously.
 - Then draw from these probability distributions and plot the results.



- Discuss your findings.

Discrete Fourier sine transform:

The square integrable, twice differentiable functions on an interval $[0, L]$, which vanish at the boundaries are spanning a Hilbert space \mathcal{F} .

An ON basis of \mathcal{F} is given by the set of functions
Ex "Hilbert basis"

$$b_n(x) := \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right), \text{ where } n = 1, 2, \dots$$

To get you started: On Minkowski space

Consider the Klein Gordon equation

$$(\partial_t^2 - \Delta + m^2)\phi(x, t) = 0$$

in a box $[0, L] \times [0, L] \times [0, L]$ with Dirichlet boundary conditions:

$$\phi(\text{boundary}) = 0$$

Recall that if the box is chosen large enough, the physics of the boundaries does not matter in the middle.

But the above boundary conditions have the mathematical advantage that we can use the discrete Fourier sine transform:

Tasks: * Use this transform to obtain a mode decomposition of $\phi(x, t)$ with coefficients $\hat{\phi}_m(t)$.

* Quantize by translating the equations

$$(\partial_t^2 - \Delta + m^2)\hat{\phi}(x, t) = 0$$


$$[\hat{\phi}(x, t), \hat{\phi}(x', t)] = i \delta(x - x') \quad \text{etc}$$

$$\hat{\phi}(x, t)^\dagger = \hat{\phi}(x, t)$$

(m_1, m_2, m_3)

calculate the probability distribution for finding certain outcomes when 'measuring' all $\hat{\phi}(x)$ simultaneously.

- Then draw from these probability distributions and plot the results.



- Discuss your findings.

in a box $[0, L] \times [0, L] \times [0, L]$ with Dirichlet boundary conditions:

$$\phi(\text{boundary}) = 0$$

Recall that if the box is chosen large enough, the physics of the boundaries does not matter in the middle.

But the above boundary conditions have the mathematical advantage that we can use the discrete Fourier sine transform:

Discrete Fourier sine transform:

The square integrable, twice differentiable functions on an interval $[0, L]$, which vanish at the boundaries are spanning a Hilbert space \mathcal{F} .

An ON basis of \mathcal{F} is given by the set of functions
Ex "Hilbert basis"

$$b_n(x) := \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right), \text{ where } n = 1, 2, \dots$$

i.e., we have $\int_0^L b_n(x) b_m(x) dx = \delta_{nm}$ and therefore for any $f \in \mathcal{F}$:

$$f(x) = \sum_{n=1}^{\infty} f_n \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right) \text{ with } f_n := \int_0^L f(x) \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right) dx.$$

Tasks: * Use this transform to obtain a mode decomposition of $\hat{\phi}(x, t)$ with coefficients $\hat{\phi}_m(t)$.

(m_1, m_2, m_3)

* Quantize by translating the equations

$$(\partial_t^2 - \Delta + m^2) \hat{\phi}(x, t) = 0$$

$$[\hat{\phi}(x, t), \hat{\phi}(x', t)] = i \delta(x - x') \quad \text{etc}$$

$$\hat{\phi}(x, t)^* = \hat{\phi}(x, t)$$

into equations that the $\hat{\phi}_m(t)$ must obey.

* You should arrive at harmonic oscillators. Assume they are in their joint ground state, which is the vacuum state. Calculate the probability distribution for each $\hat{\phi}_m(t)$

An **ON basis** of \mathcal{F} is given by the set of functions
↳ a "Hilbert basis"

$$b_n(x) := \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right), \text{ where } n = 1, 2, \dots$$

i.e., we have $\int_0^L b_n(x) b_m(x) dx = \delta_{nm}$ and therefore for any $f \in \mathcal{F}$:

$$f(x) = \sum_{n=1}^{\infty} f_n \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right) \text{ with } f_n := \int_0^L f(x) \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right) dx.$$

* Draw ϕ_n measurement outcomes from those distributions and plot the resulting $\phi(x)$.

- In practice, you can only use finitely many coefficients ϕ_n . How does the resulting picture change as you take more and more coefficients into account? What do you expect in the limit of all ϕ_n taken into account?
- What effect does the mass m have?
- Plot a case when space has 2 dimensions and compare with 2-dim slices of cases of space having more dimensions. (In each case, keep the longest w in any direction the same.)
- Draw and plot the case of a wave packet state.

$$(\partial_t^2 - \Delta + m^2) \phi(x, t) = 0$$

$$[\hat{\phi}(x, t), \hat{\phi}(x', t)] = i \delta(x - x') \quad \text{etc}$$

$$\hat{\phi}(x, t)^\dagger = \hat{\phi}(x, t)$$

into equations that the $\hat{\phi}_n(t)$ must obey.

* You should arrive at harmonic oscillators. Assume they are in their joint ground state, which is the vacuum state. Calculate the probability distribution for each $\hat{\phi}_n(t)$.


Format of project report:

- 15-20 pages
- Title / Abstract / Introduction / Motivation / Theory / Method / Results / Discussion
 : repeated for each sub-project
- Conclusions / Suggestions for further study
- Bibliography and software used
- No need to stick to exactly that format.

Recall: Descriptions are fine but explanations are what we are after. 13/6

QFT for Cosmology, Achim Kempf, Lecture 11

Recall the strategy:



Curvature:

- We postulate the coordinate system-independent Klein Gordon action:

$$S[\phi] = \frac{1}{2} \int_{\Sigma} (g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2 - V(\phi, t)) \sqrt{|g|} d^4x$$

- We will allow almost arbitrary metric tensors $g_{\mu\nu}(x)$, even those for which there do not exist coordinates \tilde{x} in which:

$$\tilde{g}_{\mu\nu}(\tilde{x}) = \eta_{\mu\nu} \text{ for all } \tilde{x}$$

- But we must have that, at least locally, special relativity holds!

⇒ Consider only $g_{\mu\nu}(x)$ for which for each x_0 , there exists a change of coordinates

$$x \rightarrow \tilde{x}$$

so that:

$$\tilde{g}_{\mu\nu}(\tilde{x}_0) = \eta_{\mu\nu}$$

□ This requirement is The Equivalence Principle:

- * We postulate that gravity can always locally be eliminated:
- * We assume that if a freely falling observer in a small region sets up a rectangular coordinate system the observer will see arbitrarily small gravity effects if the region is made arbitrarily small.

How can one identify the presence of curvature?


- * Assume we are given a metric tensor $g_{\mu\nu}(x)$ as an explicit matrix-valued function, in some coordinates.
- * How can we determine whether or not this is, e.g., the metric tensor of flat space-time, i.e., whether or not there exist coordinates \tilde{x} in which $\tilde{g}_{\mu\nu}(\tilde{x}) = \eta_{\mu\nu}$?

- * We assume that if a freely falling observer in a small region sets up a rectangular coordinate system the observer will see arbitrarily small gravity effects if the region is made arbitrarily small.
- * For this to be true, any body's gravitational mass must be equal to its inertial mass, i.e. all bodies must fall equally. (Else the notion of freely falling observer is not even well defined)

Define: The "Riemann Curvature Tensor":

$$R^i{}_{jkl}(x) := \Gamma^i_{ljk}(x) - \Gamma^i_{kjl}(x) + \Gamma^p_{lj}(x)\Gamma^i_{kp}(x) - \Gamma^p_{jk}(x)\Gamma^i_{lp}(x)$$

It's rôle? A space is called curved at x if the parallel transport of a vector v along an infinitesimal parallelogram returns the vector v' to x , but v' is rotated by some amount. $R^i{}_{jkl}$ tells by how much:



$$(v' - v)^a = \eta^a{}_b R^b{}_{cd}(x) v^c v^d$$

- * How can we determine whether or not this is, e.g., the metric tensor of flat space-time, i.e., whether or not there exist coordinates \tilde{x} in which $\tilde{g}_{\mu\nu} = \eta_{\mu\nu}$?
- * This problem is solved in differential geometry:

Define: The "Christoffel symbol functions":

$$\Gamma^{\alpha}{}_{\beta\gamma}(x) := \frac{1}{2} g^{\alpha\delta}(x) (g_{\delta\gamma,\beta}(x) + g_{\beta\delta,\gamma}(x) - g_{\beta\gamma,\delta}(x))$$

Remark:

If the parallelogram does not even close we say that space-time has "Torsion". There is no evidence for torsion in nature.

Proposition:


Assume that, in a region, A , of space-time:

$$R^i{}_{jkl}(x) = 0 \text{ for all } x \in A$$

Then and only then there exist coordinates \tilde{x} so that:

$$\tilde{g}_{\mu\nu}(\tilde{x}) = \eta_{\mu\nu} \text{ for all } x \in A$$

transport of a vector v along an infinitesimal parallelogram returns the vector v' to x , but v' is rotated by some amount. $R^{\mu}_{\nu\sigma\epsilon}$ tells by how much:



$$(v' - v)^{\alpha} = \gamma^{\alpha}_{\nu} R^{\alpha}_{\nu\sigma\epsilon}(x) v^{\nu}$$

Proposition:

Assume that, in a region, A , of space-time:

$$R^{\mu}_{\nu\sigma\epsilon}(x) = 0 \text{ for all } x \in A$$

Then and only then there exist coordinates \tilde{x} so that:

$$\tilde{g}_{\mu\nu}(\tilde{x}) = \eta_{\mu\nu} \text{ for all } x \in A$$

The dynamics of space-time

Problem: What are the equations of motion for spacetime's curvature?

Which are the degrees of freedom of curvature for which we have to find an equation of motion?

- We saw that the curvature of space-time is encoded in the matrix-valued metric function:

$$g_{\mu\nu}(x)$$

- However, if $g_{\mu\nu}(x)$ looks nontrivial, this can be for two different reasons:

1. Spacetime has little or no curvature and $g_{\mu\nu}(x)$ is nontrivial just because of an unlucky choice of coordinates.

2. Space time is curved, i.e., we can not make $g_{\mu\nu}(x)$ take the form $\tilde{g}_{\mu\nu}(\tilde{x}) = \eta_{\mu\nu}$ for all \tilde{x} no matter which coordinates we choose.

□ Therefore, it is difficult to pinpoint in the matrix function $g_{\mu\nu}(x)$ the curvature degrees of freedom.

□ And: Even the entries $R^{\mu}_{\nu\sigma\epsilon}$ of the curvature tensor are coordinate system dependent.

- We saw that the curvature of space-time is encoded in the matrix-valued metric function: $g_{\mu\nu}(x)$
- However, if $g_{\mu\nu}(x)$ looks non-trivial, this can be for two different reasons:

Strategy:

- Use the degrees of freedom of curvature to build a scalar and therefore coordinate system independent function $S_{\mu\nu}[g]$
- Then, use this function as the action for gravity.
- The equations of motion for gravity should follow from the action principle (and they do).

→ Need to define a scalar that encodes curvature!
We begin by going from a 4-tensor to a 2-tensor:

no matter which coordinates we choose.

- Therefore, it is difficult to pinpoint in the metric function $g_{\mu\nu}(x)$ the curvature degrees of freedom.
- And: Even the entries $R^{\mu}_{\nu\sigma\lambda}$ of the curvature tensor are coordinate system dependent.

Definition: The "Ricci tensor"

$$R_{\mu\nu}(x) := R^{\lambda}_{\mu\lambda\nu}(x)$$

Recall: $\sum_{\lambda=0}^3$ is implied

Note: Other index contractions would vanish because of antisymmetries of $R^{\mu}_{\nu\sigma\lambda}(x)$ that are implied by the definition of $R^{\mu}_{\nu\sigma\lambda}(x)$.

Remark:

$R_{\mu\nu}(x)$ carries strictly less information than the full Riemann curvature tensor:

- * If $R_{\mu\nu}(x) = 0$ it is still possible that $R^{\mu}_{\nu\sigma\lambda}(x) \neq 0!$
- * This happens to be the case, e.g., for gravitational waves.

Then, use this function as the action for gravity.

The equations of motion for gravity should follow from the action principle (and they do).

Need to define a scalar that encodes curvature!
We begin by going from a 4-tensor to a 2-tensor:

of antisymmetries of $R^{\mu\nu\alpha\beta}(x)$ that are implied by the definition of $R^{\mu\nu\alpha\beta}(x)$.

Remark:

$R_{\mu\nu}(x)$ carries strictly less information than the full Riemann curvature tensor:

- * If $R_{\mu\nu}(x) = 0$ it is still possible that $R^{\mu\nu\alpha\beta}(x) \neq 0$!
- * This happens to be the case, e.g., for gravitational waves.

Definition: The "curvature scalar" (or "Ricci scalar")

$$R(x) := g^{\mu\nu}(x) R_{\mu\nu}(x)$$

Other curvature scalars:

- * The simplest scalar that can be formed from the metric alone is $g^{\mu\nu}(x) g_{\mu\nu}(x) = 4$.
- * The next simplest scalar that can be formed is the Ricci scalar $R(x)$.
- * All other scalars made out of g only are composed of higher powers of the Riemann tensor $R^{\mu\nu\alpha\beta}(x)$:

$$R_{\mu\nu} R^{\mu\nu}, R_{\mu\nu} R^{\nu\alpha} R_{\alpha\mu}, \text{ etc.}$$

The gravitational action

A priori, the full action now reads:

$$S_{\text{tot}}[g, \phi, e_i] = S_{\text{gc}} + S_{\text{matter}} + S_{\text{grav}}$$

other "matter" fields for e, quads, photons etc

Euler-Lagrange action

with: $S_{\text{grav}}[g] := \int (c_0 + c_1 R(x) + c_2 R_{\mu\nu} R^{\mu\nu}(x) + c_3 R_{\mu\nu} R^{\nu\alpha} R_{\alpha\mu} + \dots) \sqrt{|g|} d^4x$

Comparison with experiment shows evidence only for the first two terms:

$$S_{\text{grav}}[g] = -\frac{1}{16\pi G} \int (2\Lambda + R(x)) \sqrt{|g|} d^4x$$

"Einstein action"
Newton's constant
"Cosmological constant"

Remark: D. Lovelock here determined all generalizations to higher terms and higher dimensions that still possess 2nd order initial value problem Lat Applied Math at UW

- * The next simplest scalar that can be formed is the Ricci scalar $R(x)$.
- * All other scalars made out of g only are composed of higher powers of the Riemann tensor $R^{\mu\nu\sigma\lambda}(x)$:
 $R_{\mu\nu}R^{\mu\nu}$, $R_{\mu\nu}R^{\nu\sigma}R_{\sigma\mu}$ etc.

with: $\int \sqrt{|g|} d^4x$

Comparison with experiment shows evidence only for the first two terms:

$$S_{\text{grav}}[g] = -\frac{1}{16\pi G} \int (2\Lambda + R(x)) \sqrt{|g(x)|} d^4x$$

"Einstein action"
Newton's constant
"Cosmological constant"

Remark: D. Lovelock here determined all generalizations to higher terms and higher dimensions that still possess 2nd order initial value problems.

The equations of motion

The action principle is to require that the action be extremal with respect to all degrees of freedom:

A) $\frac{\delta S_{\text{tot}}}{\delta \ell_i(x)} = 0$ B) $\frac{\delta S_{\text{tot}}}{\delta g_{\mu\nu}(x)} = 0$ C) $\frac{\delta S_{\text{tot}}}{\delta \phi(x)} = 0$

A) Require: $\frac{\delta S_{\text{tot}}[g, \ell, \phi]}{\delta \ell_i(x)} = 0$

This yields the general relativistically covariant field equations for all "other" fields. (We will ignore the $\ell_i(x)$ for now.)

Quantization: Legendre transform $\rightarrow H(\ell_i, \pi_{\ell_i}) \rightarrow$ impose $\{\ell_i, \pi_{\ell_i}\} = \delta$ etc.

* Quantization: To quantize the Einstein equation

B) Require: $\frac{\delta}{\delta g_{\mu\nu}(x)} S_{\text{tot}}[\phi, \ell, g] = 0$

This yields the equation of motion for the dynamics of curvature, i.e., the Einstein equation:

(See exercise in Mukhanov's text) \rightarrow $R_{\mu\nu}(x) - \frac{1}{2}g_{\mu\nu}(x)R(x) + \Lambda g_{\mu\nu}(x) = +8\pi G T_{\mu\nu}(x)$

$\sim \frac{\delta S_{\text{grav}}}{\delta g_{\mu\nu}(x)}$
 $\sim \frac{\delta(S_{\text{matter}} + S_{\text{cosmo}})}{\delta g_{\mu\nu}(x)}$

* Here, $T_{\mu\nu}(x)$ is the "Energy Momentum Tensor". Neglecting the contribution by the $\ell_i(x)$, one obtains:

$$T_{\mu\nu}^{(K+\phi)} = \phi_{,\mu}(x)\phi_{,\nu}(x) - g_{\mu\nu}(x) \left(\frac{1}{2}g^{\alpha\beta}(x)\phi_{,\alpha}(x)\phi_{,\beta}(x) - V(\phi(x)) \right)$$

mass term in ϕ included

C) Require: $\frac{\delta}{\delta \phi(x)} S_{\text{tot}}[g, \phi] = 0$

A) Require: $\frac{\delta S_{\text{tot}}[g, \ell, \phi]}{\delta \ell(x)} = 0$

This yields the general relativistically covariant field equations for all "other" fields. (We will ignore the $\ell_i(x)$ for now.)

Quantization: Legendre transform $\rightarrow H(\ell, \Pi_\ell) \rightarrow$ impose $\{\ell, \Pi_\ell\} = \delta$ etc.

* Quantization: To quantize the Einstein equation is difficult for many reasons:

- o For example, it is difficult to separate the curvature degrees of freedom from mere artifacts of the choice of the coordinate system.
- o Also, the Einstein equation is highly nonlinear.
- o So far, all attempts have run into severe difficulties, even perturbative approaches.

- \rightsquigarrow This course:
- 1.) We will have initially consider known classical solutions $g_{\mu\nu}(x)$ and quantize only $\phi(x)$.
 - 2.) Then, we will quantize linear perturbations of the metric.

$$\sim \frac{\delta S_{\text{grav}}}{\delta g_{\mu\nu}(x)} \quad \sim - \frac{\delta(S_{\text{matter}} + S_{\text{cosmo}})}{\delta g_{\mu\nu}(x)}$$

* Here, $T_{\mu\nu}(x)$ is the "Energy Momentum Tensor". Neglecting the contribution by the $\ell_i(x)$, one obtains:

$$T_{\mu\nu}^{(k+\ell)}(x) = \phi_{,\mu}(x)\phi_{,\nu}(x) - g_{\mu\nu}(x) \left(\frac{1}{2} g^{\alpha\beta}(x) \phi_{,\alpha}(x)\phi_{,\beta}(x) - V(\phi(x)) \right)$$

mass term $m^2 \phi^2$ included

C) Require: $\frac{\delta}{\delta \phi(x)} S_{\text{tot}}[g, \phi] = 0$

□ Since ϕ occurs only in S_{tot} we have, equivalently:

$$\frac{\delta S_{\text{tot}}}{\delta \phi(x)} = 0$$

□ Recall S_{tot} :

$$S_{\text{tot}}[\phi] = \frac{1}{2} \int_{\mathcal{M}} \left(g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - \underbrace{m^2 \phi^2 - \lambda \phi^4}_{\text{Example of a potential}} \right) \sqrt{|g|} d^4x$$

□ Apply the Euler Lagrange equations:

$$\frac{\delta S_{\text{tot}}[\phi, \{g\}, g]}{\delta \phi(x, t)} = \partial_\mu \frac{\delta S[\phi, \{g\}, g]}{\delta (\phi_{,\mu}(x, t))}$$

- Also, the Einstein equation is highly nonlinear.
- So far, all attempts have run into severe difficulties, even perturbative approaches.

→ This course: 1.) We will here initially consider known classical solutions $g_{\mu\nu}(x)$ and quantize only $\phi(x)$.
 2.) Then, we will quantize linear perturbations of the metric.

⇒ Klein Gordon equation in general relativity:

$$\left(-\frac{1}{2} m^2 \square \phi(x) - \frac{1}{2} \lambda \phi^3(x)\right) \sqrt{|g(x)|} = \partial_\mu \left(\frac{\partial}{\partial x^\mu} \phi(x) \sqrt{|g(x)|}\right)$$

Thus:

$$\frac{1}{\sqrt{|g(x)|}} \frac{\partial}{\partial x^\mu} \left(g^{\mu\nu}(x) \sqrt{|g(x)|} \phi_{,\nu}(x) \right) + m^2 \phi(x) + 2\lambda \phi^3(x) = 0$$

Definition: The "d'Alembert operator", \square

$$\square := \frac{1}{\sqrt{|g(x)|}} \frac{\partial}{\partial x^\mu} g^{\mu\nu}(x) \sqrt{|g(x)|} \frac{\partial}{\partial x^\nu}$$

Thus:

$$\square \phi(x) + m^2 \phi(x) + 2\lambda \phi^3(x) = 0$$

Recall S_{KG} :


$$S_{KG}[\phi] = \frac{1}{2} \int_{\mathcal{M}} \left(g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2 - \lambda \phi^4 \right) \sqrt{|g|} d^4x$$

Example of a potential

Apply the Euler Lagrange equations:

$$\frac{\delta S_{KG}[\phi; \{g\}]}{\delta \phi(x,t)} = \partial_\mu \frac{\delta S_{KG}[\phi; \{g\}]}{\delta (\phi_{,\mu}(x,t))}$$

Next: Step 3 in



□ ⇒ Klein Gordon equation in general relativity:

$$\left(-\frac{1}{2} m^2 \Delta \phi(x) - \frac{1}{2} \lambda \phi^3(x)\right) \sqrt{|g(x)|} = \partial_\mu \left(\frac{\partial}{\partial x^\mu} \phi(x) \sqrt{|g(x)|} \right)$$

Thus:

$$\frac{1}{\sqrt{|g(x)|}} \frac{\partial}{\partial x^\mu} \left(g^{\mu\nu}(x) \sqrt{|g(x)|} \phi_{,\nu}(x) \right) + m^2 \phi(x) + \lambda \phi^3(x) = 0$$

□ Definition: The "d'Alembert operator", □

$$\square := \frac{1}{\sqrt{|g(x)|}} \frac{\partial}{\partial x^\mu} g^{\mu\nu}(x) \sqrt{|g(x)|} \frac{\partial}{\partial x^\nu}$$

Thus:

$$\square \phi(x) + m^2 \phi(x) + \lambda \phi^3(x) = 0$$

Next: Step 3 in

SR, 1st Q
Hamiltonian formalism

step 1
Legendre transform
(equivalence) →

SR, 1st Q.
Lagrangian formalism

step 2 ↓ allow curvature

GR, 1st Q
Hamiltonian formalism

step 3
Legendre transform
(equivalence) ←

GR, 1st Q
Lagrangian formalism

step 4 ↓
GR, 2nd Q
Hamiltonian formalism

Dirac-Schwinger eqns are same
(equivalence) ←

(PI) ↓
GR, 2nd Q
Lagrangian formalism
(Path integral QFT)



Comment on step (PI): 2nd quantization with path integral

□ Assume a fixed spacetime is chosen and we are given its metric $g_{\mu\nu}(x)$ in some arbitrary coordinate system.

□ Then, for each field $\phi(\vec{x}, t)$ we can calculate its action $S_{int}[\phi, g]$:

$$S_{int}[\phi, g] = \frac{1}{2} \int_{\Sigma} \left(g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2 - \lambda \phi^3 \right) \sqrt{|g|} d^4x$$

□ Consider, e.g., the vacuum expectation value of $\phi(\vec{x}, t) \phi(\vec{x}', t)$, i.e., the correlation function of field amplitudes:

$$G(\vec{x}, t, \vec{x}', t) = \langle 0 | \hat{\phi}(\vec{x}, t) \hat{\phi}(\vec{x}', t) | 0 \rangle$$

□ We will later see how to calculate it using commutation relations etc.

□ With Feynman we also get it from the path integral:

Then, for each field $\phi(x, t)$ we can calculate

action $S_{KG}[\phi, g]$:

$$S_{KG}[\phi, g] = \frac{1}{2} \int_{\mathbb{R}^4} (g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2) \sqrt{|g|} d^4x$$

Following Feynman, we obtain 'probability amplitudes':

$$\text{prob. ampl. } [\phi] := e^{\frac{i}{\hbar} S_{KG}[\phi, g]}$$

We will later see how to calculate it using commutation relations etc.

With Feynman we also get it from the path integral:

Advantages:

- 1) Quick derivation of Feynman rules
- 2) Manifestly covariant


Problems:

- 1) Not defined due to uncountable number of configurations and related divergences.
- 2) Even when these are temporarily regularised, the identification of the measure still is ambiguous.
 → This issue is better handled in canonical formalism.

$$G(\vec{x}, t; \vec{x}', t') = N \int_{\phi(\vec{x}', t')}^{\phi(\vec{x}, t)} e^{\frac{i}{\hbar} S_{KG}[\phi, g]} \mathcal{D}[\phi]$$

↑ Path Integral "over a function space"

Recall strategy:



Recall:

General relativistic covariant Klein Gordon theory in the Lagrangian formulation (neglecting the potential):

$$S_{KG} = \frac{1}{2} \int_{\mathbb{R}^4} (g^{\mu\nu}(x) \phi_{,\mu}(x) \phi_{,\nu}(x) - m^2 \phi^2(x)) \sqrt{|g|} d^4x$$

we assume that the coordinate system is such, for simplicity.

It yields, via $\frac{\delta S_{KG}}{\delta \phi} = 0$ the Klein Gordon eqn:

$$\frac{1}{\sqrt{|g|}} \frac{\partial}{\partial x^\mu} (g^{\mu\nu}(x) \sqrt{|g(x)|} \phi_{,\nu}(x)) + m^2 \phi(x) = 0 \quad (KG)$$

formalism

GR, 1st Q
Hamiltonian
formalism

step 4

GR, 2nd Q
Hamiltonian
formalism

step 3
Legendre transform
(equivalence)

Dyson Schwinger eqns are same
(equivalence)

formalism

step 2 allow curvature

GR, 1st Q
Lagrangian
formalism

we are here

GR, 2nd Q
Lagrangian formalism
(Path integral $\int \mathcal{D}\phi$)

\mathbb{R}^4 we assume that the coordinate system is such, for simplicity.

It yields, via $\frac{\delta S_{KG}}{\delta \phi} = 0$ the Klein Gordon eqn:

$$\frac{1}{\sqrt{|g|}} \frac{\partial}{\partial x^\mu} \left(g^{\mu\nu} \sqrt{|g|} \partial_\nu \phi \right) + m^2 \phi(x) = 0 \quad (KG)$$

* We read off the Lagrangian:

$$L_{KG}(\phi) = \frac{1}{2} \int_{\mathbb{R}^3} \left(g^{\mu\nu}(x,t) \phi_{,\mu}(x,t) \phi_{,\nu}(x,t) - m^2 \phi^2(x,t) \right) \sqrt{|g|} d^3x$$

Step 3: Legendre transform back to the Hamiltonian form

* The transform:

$$H(\phi, \pi, t) \xleftarrow[\text{Legendre transform}]{\pi(x,t) := \frac{\delta L}{\delta \phi_{,\mu}(x,t)}} L(\phi, \phi_{,\mu}, t)$$

* Thus, the canonically conjugate field $\pi(x,t)$ reads:

$$\pi(x,t) = \frac{\delta L}{\delta \phi_{,\mu}(x,t)} = \sqrt{|g(x,t)|} g^{0\mu}(x,t) \phi_{,\mu}(x,t)$$

* Explicitly:

$$\pi(x,t) = \sqrt{|g|} g^{00} \phi_{,0} + \sum_{i=1}^3 \sqrt{|g|} g^{0i} \phi_{,i}$$

* Thus, we can also express $\phi_{,i}(x,t)$ in terms of $\phi(x,t)$ and $\pi(x,t)$ (as will be necessary after the Legendre transform):

$$\phi_{,i}(x,t) = \frac{\pi(x,t)}{g^{00} \sqrt{|g|}} - \sum_{j=1}^3 \frac{g^{0j}}{g^{00}} \phi_{,j}(x,t) \quad (V)$$

(of course, g^{00} depends on x, t too)

* The Hamiltonian:

$$H(\phi, \pi) = \int_{\mathbb{R}^3} \pi(x,t) \phi_{,0}(x,t) d^3x - \frac{1}{2} \int_{\mathbb{R}^3} \left(g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} - m^2 \phi^2 \right) \sqrt{|g|} d^3x$$

↑
Why don't we need a factor of $\sqrt{|g|}$ for covariance here? Because π has it built in!

$= L(\phi, \phi_{,\mu}(\phi, \pi), t)$

Step 3: Legendre transform back to the Hamiltonian form

* The transform:

$$H(\phi, \pi, t) \xleftarrow[\text{Legendre transform}]{\pi(x,t) := \frac{\delta L}{\delta \dot{\phi}_\mu(x,t)}} L(\phi, \dot{\phi}_\mu, t)$$

* Thus, the canonically conjugate field $\pi(x,t)$ reads:

$$\pi(x,t) = \frac{\delta L}{\delta \dot{\phi}_\mu(x,t)} = \sqrt{|g(x,t)|} g^{0\mu}(x,t) \dot{\phi}_{,\mu}(x,t)$$

* In H , one needs to express all occurring $\dot{\phi}_\mu$ in terms of the new variables ϕ and π , by using (V), to obtain $H(\phi, \pi, t)$.

→ Exercise: Calculate $H(\phi, \pi, t)$ and simplify the expression as far as possible.

* The equations of motion:

We know from the general properties of the Legendre transform that the equations of motion now take the form:

and $\pi(x,t)$ (as will be necessary after the Legendre transform):

$$\phi_{,\nu}(x,t) = \frac{\pi(x,t)}{g^{0\nu} \sqrt{|g|}} - \sum_{i=1}^3 \frac{g^{0i}}{g^{0\nu}} \dot{\phi}_{,i}(x,t) \quad (V)$$

(of course, $g^{0\nu}$ depends on x, t too)

* The Hamiltonian:

$$H(\phi, \pi) = \int_{R^3} \pi(x,t) \dot{\phi}_{,\nu}(x,t) d^3x - \frac{1}{2} \int_{R^3} (g^{\mu\nu} \dot{\phi}_{,\mu} \dot{\phi}_{,\nu} - m^2 \phi^2) \sqrt{|g|} d^3x$$


↑
Why don't we need a factor of $\sqrt{|g|}$ for covariances here? Because π has it built in!

= $L(\phi, \dot{\phi}_\mu, \phi, \pi, t)$

$$\frac{d}{dt} \phi(x,t) = \frac{\delta H(\phi, \pi, t)}{\delta \pi(x,t)}, \quad \frac{d}{dt} \pi(x,t) = - \frac{\delta H(\phi, \pi, t)}{\delta \phi(x,t)}$$

* Exercise: Verify that these eqns are equivalent to (KG).

We are now ready to 2nd quantize:



$$H(\phi, \pi, t) \xleftarrow[\text{Legendre transform}]{\pi(x,t) := \frac{\partial L}{\partial \dot{\phi}_\mu(x,t)}} L(\phi, \dot{\phi}_\mu, t)$$

* Thus, the canonically conjugate field $\pi(x,t)$ reads:

$$\pi(x,t) = \frac{\delta L}{\delta \dot{\phi}_\mu(x,t)} = \sqrt{|g(x,t)|} g^{\mu\nu}(x,t) \dot{\phi}_\nu(x,t)$$

* In H , one needs to express all occurring $\dot{\phi}_\mu$ in terms of the new variables ϕ and π , by using (V), to obtain $H(\phi, \pi, t)$.

→ Exercise: Calculate $H(\phi, \pi, t)$ and simplify the expression as far as possible.

* The equations of motion:

We know from the general properties of the Legendre transform that the equations of motion now take the form:

* The Hamiltonian:

$$H(\phi, \pi) = \int_{\mathbb{R}^3} \pi(x,t) \dot{\phi}_\mu(x,t) d^3x - \frac{1}{2} \int_{\mathbb{R}^3} (g^{\mu\nu} \dot{\phi}_\mu \dot{\phi}_\nu - m^2 \phi^2) \sqrt{|g|} d^3x$$

↑
Why don't we need a factor of $\sqrt{|g|}$ for covariance here? Because π has it built in!

$= L(\phi, \dot{\phi}_\mu(\phi, \pi), t)$

$$\frac{d}{dt} \phi(x,t) = \frac{\delta H(\phi, \pi, t)}{\delta \pi(x,t)}, \quad \frac{d}{dt} \pi(x,t) = - \frac{\delta H(\phi, \pi, t)}{\delta \phi(x,t)}$$

* Exercise: Verify that these eqns are equivalent to (K6).

We are now ready to 2nd quantize:

