Title: Twisted Tools for (Untwisted) Quantum Field Theory

Speakers: Justin Kulp

Collection: Higher Categorical Tools for Quantum Phases of Matter

Date: March 18, 2024 - 2:00 PM

URL: https://pirsa.org/24030080

Twisted Tools for (Untwisted) Quantum Field Theory

Higher Categorical Tools for Quantum Phases of Matter

Justin Kulp
with Davide Gaiotto, and Jingxiang Wu
and earlier works with Kasia Budzik,
Brian Williams, and Matthew Yu.

Simons Center for Geometry and Physics

18/Mar/2024

arXiv:2403.tomorrow, arXiv:2306.01039, arXiv:2207.14321

Outlines and Punchlines 1/2

- Discuss ideas from formal deformation theory in QFT
  - Familiar example is Ocneanu rigidity of fusion categories
- QFTs have "higher" **multilinear k-ary operations** ("brackets")

$$\{-,-,\ldots,-\} \tag{1}$$

  - Control: **deformations**, (generalized) **OPEs**, and **anomalies**
  - ∞-algebras, factorization algebras, and operads
- Familiar to high energy physicists and mathematicians who have studied twisted SQFTs (descent relations)
  - Not limited to twisted scenarios
- Can go very far in the case of (mixed) Holomorphic and/or Topological (HT) theories

Outlines and Punchlines 2/2

I. Reminder on beta-function and BRST-symmetry
II. Introduce the eta-function and higher brackets
III. Example of computing the eta-function
IV. Generalizations to other scenarios
V. Quick survey of some results in holomorphic-topological scenarios

Three Punchlines

1. η-vector exists and is computable
2. η-vector contains anomalies, OPEs, and more
3. Non-renormalization theorem for HT theories

Deformations of QFTs

- Given a QFT T, it can be deformed by turning on interactions

$$S_T + \sum_i g^i \int_{\mathbb{R}^d} \mathcal{O}^i(x) d^d x \qquad (2)$$



  - g^i are coordinates on theory space
  - Work perturbatively in couplings g^i
- Defines a **formal pointed neighbourhood** $\mathcal{D}[T]$ of T, consisting of all effective QFTs obtained by perturbative deformation of T
  - Pointed because there is a distinguished point, called T.
  - Formal because we only consider deformations in an infinitesimal nbd of T (we are not at finite coupling).
  - Think of formal/infinitesimal as synonym for "perturbative"

The Beta Function

- Generic QFT (point) is not scale invariant
  - Scale transformation on T is traded for a change of the couplings e.g. integrating out DoF modifies couplings
- We encode infinitesimal scale transformations in vector field on theory space, the **beta function**

$$\boldsymbol{\beta} = \sum_i \beta^i(g) \frac{\partial}{\partial g^i} \,. \tag{3}$$



  - Perturb around (typically free) scale-invariant theory, $\boldsymbol{\beta} = 0$
  - Deformations of T preserving scale invariance are zeroes of $\boldsymbol{\beta}$
- The coefficients $\beta^i(g)$ are power series in g

$$\beta^i(g) = \underbrace{(d - \Delta_i)}_{\text{Classical}} g^i + O(g^2) \tag{4}$$

  - Tune relevant terms to 0 and study $\boldsymbol{\beta}$ as a measure of scale generated by "quantum effects"

The BV-BRST Formalism 1/2

- Quantization of non-abelian gauge theories is hard: formulated redundantly in exchange for other properties

$$Z = \int [DAD\bar{\psi}D\psi] e^{-S[A,\bar{\psi},\psi]} =: \int [D\Phi] e^{-S[\Phi]}\,, \tag{5}$$

  - Introduce a gauge fixing procedure and Fadeev-Poppov ghosts b and c

$$Z = \int [D\Phi DB_A db_A dc^\alpha] e^{-S[\Phi] + iB_A F^A[\Phi] - b_A c^\alpha \delta_\alpha F^A[\Phi]} \tag{6}$$

- Gauge fixed action still has residual nilpotent odd global symmetry involving fields and ghosts, called **BRST symmetry**.

$$\begin{aligned} \delta_{\mathrm{BRST}}\Phi &= -i\epsilon c^\alpha \delta_\alpha \Phi\,, & \delta_{\mathrm{BRST}} B_A &= 0\,, \\ \delta_{\mathrm{BRST}} c^\alpha &= \frac{i}{2}\epsilon f^\alpha_{\beta\gamma} c^\beta c^\gamma\,, & \delta_{\mathrm{BRST}} b_A &= \epsilon B_A\,, \end{aligned} \tag{7}$$

  - Physical theory can be identified with Q_{BRST}-cohomology

The BV-BRST Formalism 2/2

- Focus on theories defined in BV-BRST formalism:
  - T is embedded in a bigger ambient theory $\widetilde{T}$ with ghosts, anti-ghosts, anti-fields, etc.
  - Grassmann odd nilpotent symmetry Q_{BRST}
  - Observables in T are recovered from $\widetilde{T}$ by taking Q_{BRST} coho

$$\begin{aligned} \mathrm{Ops}_T &= (\mathrm{Ops}_{\widetilde{T}}, Q_{\mathrm{BRST}}) \\ \mathrm{Int}_T &= (\mathrm{Ops}_{\widetilde{T}}[dx], \mathrm{d} + Q_{\mathrm{BRST}}) \end{aligned} \tag{8}$$

- i.e. we will work in BV formalism
  - Essential to quantizing p-form gauge theories, theories which only close on-shell, field-dependent structure constants, or theories with other complicated constraints
  - Not restricted to such complicated theories either

The Eta Function

- Can compute analog of β **for any type of transformation**.
  - Ex. Non-relativistic scale transformations $(t, x) \mapsto (\lambda^z t, x)$
  - Ex. Anomalous axial transformation on θ angle in gauge theory
- Consider $T \hookrightarrow (\widetilde{T}, Q_{\mathrm{BRST}})$ described in a BRST formalism in terms of ambient $\widetilde{T}$
  - To deform T, we deform $\widetilde{T}$ without breaking BRST symmetry
  - Consider deformations of $\widetilde{T}$ with Grassmann odd couplings, non-trivial ghost number, etc. This is formal pointed dg-supermanifold $\mathcal{D}[\widetilde{T}]$.
- BRST symmetry will be encoded in a vector field

$$\boldsymbol{\eta} = \sum_i \eta^i(g) \frac{\partial}{\partial g^i} \,. \tag{9}$$

  - Linear term tells us if adding an interaction $\mathcal{I}$ explicitly/classically violates BRST symmetry
  - Higher order terms do so "quantum mechanically"

Higher Algebra

- Since $Q^2 = 0$, the eta function $\eta^2 = 0$.
  - Wess-Zumino consistency condition for BRST symmetry
  - Gves **quadratic constraints** on coefficient functions $\eta^i(g)$

$$\eta^i(g) = \sum_{n>0} \frac{1}{n!} \sum_{j_1 \cdots j_n} \eta^i_{j_1 \cdots j_n} g^{j_1} \cdots g^{j_n} \,, \tag{10}$$

- Define the following multilinear operation $\mathrm{Int}^{\otimes n} \to \mathrm{Int}$

$$\{g^{j_1}\mathcal{I}_{j_1}, \cdots, g^{j_n}\mathcal{I}_{j_n}\} = \eta^i_{j_1 \cdots j_n} g^{j_1} \cdots g^{j_n} \mathcal{I}_i \,. \tag{11}$$

  - The BRST variation becomes

$$\boldsymbol{\eta}\,\mathcal{I} = \{\mathcal{I}\} + \frac{1}{2!}\{\mathcal{I},\mathcal{I}\} + \frac{1}{3!}\{\mathcal{I},\mathcal{I},\mathcal{I}\} + \ldots \tag{12}$$

$\boldsymbol{\eta}^2 = 0 \quad \Leftrightarrow$ The coefficients $\eta^i_{j_1 \cdots j_n}$ and brackets $\{\cdot, \ldots, \cdot\}$ define an $L_\infty[1]$**-algebra** structure on In.

Basic Eta-Function Calculation - General 1/2

- η is not just abstract fun, it is computable fun
- Correlation functions of T deformed by $\mathcal{I}$ are correlation functions of T with additional insertions

$$\langle \mathcal{O}_1(x_1) \cdots \mathcal{O}_n(x_n) \rangle_{T+\mathcal{I}} = \left\langle \mathcal{O}_1(x_1) \cdots \mathcal{O}_n(x_n) e^{g \int d^d x \, \mathcal{I}} \right\rangle_T \quad (13)$$

  - At **linear order**, the BRST anomaly generated by $\mathcal{I}_i$ is just

$$\int_{\mathbb{R}^d} [Q, \mathcal{I}_i] \,. \quad (14)$$

  - Write

$$[Q, \mathcal{I}_i] = \sum_j \eta_i^j \mathcal{I}_j + d\mathcal{J}_i \quad (15)$$

- In perturbation theory, higher order terms

$$O(g^n) \sim \int_{\mathbb{R}^{dn}} \mathcal{I}(y_1) \cdots \mathcal{I}(y_n) \quad (16)$$

  - Need **regularization** to avoid UV divergences from colliding $\mathcal{I}$

Basic Eta-Function Calculation - General 2/2

- E.g. **at** $O(g^2)$ we can regularize the deformation to

$$\int_{\mathbb{R}^{2d}} f_\epsilon^{(2)}(x_1, x_2)\mathcal{I}(x_1)\mathcal{I}(x_2) \tag{17}$$

- Now we can compute

$$\left[Q, \int_{\mathbb{R}^{2d}} f_\epsilon^{(2)}(x_1, x_2)\ \mathcal{I}(x_1)\,\mathcal{I}(x_2)\right]\Big|_{d\mathcal{J}} \tag{18}$$

$$= -\int_{\mathbb{R}^{2d}} df_\epsilon^{(2)}(x_1, x_2)(\mathcal{I}(x_1)\mathcal{J}(x_2) + \mathcal{J}(x_1)\mathcal{I}(x_2)) \tag{19}$$

- With a sharp cutoff (**point-splitting**) this becomes

$$\{\mathcal{I}, \mathcal{I}\}(x_2) \overset{\text{Sharp Cutoff}}{=} \int_{|x_{12}|=\epsilon} \mathcal{I}(x_1)\mathcal{J}(x_2) + \mathcal{J}(x_1)\mathcal{I}(x_2) \tag{20}$$

  - Anomaly appears in point-splitting regularization because total derivative terms give a boundary contribution.

Basic Eta-Function Calculation - Concrete 1/2

- 2d S_{Matter} with G global symmetry and G gauge theory

$$S_T = -\frac{1}{4}\int d^2x\, F_{\mu\nu}F^{\mu\nu} + S_{\text{Matter}} \tag{21}$$

Ex. Free fermions with vector current $J^\mu_a = \bar{\psi}\gamma^\mu t_a \psi$.

- Study the **interaction** $\mathcal{I} = A_\mu J^\mu$

- Add ghost and auxiliary fields $T \hookrightarrow (\widetilde{T}, Q)$
  - BRST transformation of $\mathcal{I}$ gives:

$$\delta_{\text{BRST}}(A_\mu J^\mu) = (\epsilon D_\mu c)J^\mu + A_\mu(i\epsilon g c J^\mu) = \epsilon\partial_\mu c J^\mu\,, \tag{22}$$

  - See $\mathcal{I}$ is BRST-closed up to total derivative $\mathcal{J} = cJ$
  - Term can potentially cause BRST anomaly

Basic Eta-Function Calculation - Concrete 2/2

- The two-bracket receives a contribution from the 2d JJ OPE:

$$\begin{aligned}\{\mathcal{I},\mathcal{I}\}(x_2) &= \int_{|x_{12}|=\epsilon} :AJ:(x_1)\ :cJ:(x_2) + :cJ:(x_1)\ :AJ:(x_2) \\ &= \oint_{S^1_{x_2}} (\,:A(x_1)c(x_2):\, + :c(x_1)A(x_2):\,)\,\langle J(x_1)J(x_2)\rangle \\ &= \#\ :c\ dA:\,(x_2)\,. \end{aligned} \tag{23}$$

  - We use $JJ \sim |x_{12}|^{-2}$, taylor expanded, and integrated by parts
  - $\#$ denotes combinatorial and rep-theoretic factors

- Recover well-known **1-loop anomaly for G-gauge theory**

$$\{A_\mu J^\mu, A_\nu J^\nu\} = \#\, cF_{12}\,. \tag{24}$$

  - In $2k$-dim, you recover anomaly from $(k+1)$-ary bracket

Recap and Further Connections

- Recap:
  - Deformations are integral to our understanding of QFT
  - Working in a BV-BRST formalism, we can introduce η that tracks violation of BRST symmetry due to interactions
  - η defines an L_∞-algebra on interactions and Maurer-Cartan equation is solved by well-defined deformations
  - η contains useful information like anomalies
- Descent operations in twisted theories
  - Higher brackets appear by colliding/integrating descendants
  - E.g. OPE and secondary product of cohomological TFTs is a 2-ary bracket
  - [Bomans, Wu] compute higher-central charges (a and c) of 4d SUSY gauge theories from brackets

Generalizations

- Can consider **position dependent interactions**: causes momentum-inflow p^i at each vertex
  - L_∞-brackets extended to $\otimes_i \mathrm{In}_{p^{(i)}} \to \mathrm{In}_{\sum_i p^{(i)}}$
  - Momentum-coloured operad

$$\{\mathcal{I}_{i_1\ p^{(1)}}\, \mathcal{I}_{i_2\ p^{(2)}}\ \cdots\ _{p^{(n-1)}}\, \mathcal{I}_{i_n}\} \tag{25}$$

- Distinguished subcase: holomorphic theories with holomorphic momentum λ recovers λ-brackets and higher n-Lie or homotopy conformal algebras
- **Auxiliary and defect systems**: the brackets of $T \times T_{\mathrm{probe}}$ extract information about T
  - Ex. 't Hooft anomaly of S_{Matter} is apparent in the non-trivial bracket when coupled to G-gauge theory T_{probe}
  - Ex. If T is topological QM, brackets of T recover Moyal commutator. Brackets with an auxiliary fermion recovers full Moyal-star product. 1d-topological defect brackets have A_∞.

Holomorphic-Topological Theories

Holomorphic-Topological Theories

- "**Holomorphic-Topological**" means flat spacetime has structure of $\mathbb{C}^H \times \mathbb{R}^T$ with coords $(x^{\mathbb{C}}, \bar{x}^{\mathbb{C}}, x^{\mathbb{R}})$
  - Anti-holomorphic translations in $\mathbb{C}^H$ and translations in $\mathbb{R}^T$ are gauge symmetries (Q_{BRST}-exact)
  - Interested in theories with action

$$\int_{\mathbb{C}^H \times \mathbb{R}^T} \left[(\Phi, \mathrm{d}\,\Phi) + \mathcal{I}(\Phi) \right] \, d^H x^{\mathbb{C}} \, . \tag{26}$$

  - Φ is a "superfield," and $dx^{\mathbb{R}}$ and $d\bar{x}^{\mathbb{C}}$ are "superspace coordinates"
- Appears in holomorphic-topological twists of SUSY theories.
  - Take cohomology of nilpotent supercharge Q_{SUSY}, i.e. "twist"
  - Translations are Q-exact; (cohomologically) HT sub-theory

$$\{Q, \bar{Q}_{\dot{\alpha}}\} = \partial_{\bar{z}^{\dot{\alpha}}} \tag{27}$$

- Can be used to prove a general non-renormalization theorem for theories with $T \geq 2$.

Holomorphic-Topological Integrals

- In such theories, we will be interested in brackets of the form

$$\{\mathcal{O}_1 \, \lambda_1 \cdots \lambda_{n-1} \, \mathcal{O}_n\} \tag{28}$$

- The **Feynman integrals** that contribute will take the form:

$$I_\Gamma(\lambda; z) = \int_{\mathcal{M}^{|\Gamma_0|-1}} \left[\prod_{v \in \Gamma_0}^{v \neq v_*} \mathrm{dVol}_v \, e^{\lambda_v \cdot x_v^{\mathbb{C}}} \right] \mathrm{d} \left[\prod_{e \in \Gamma_1} P_\epsilon(x_{e(0)} - x_{e(1)} + z_e) \right]$$

- Let's count the form degree of the integrand:
  - (Regulated) propagator P_ϵ is a $(0, H + T - 1)$ form
  - $(H + T) \times (|\Gamma_0| - 1)$ integration variables: one for each vertex of graph, and throw one vertex away by translation symmetry.
  - $(|\Gamma_1| - 1)$ regulated propagators and one $(0, H + T)$-form cut propagator $\mathrm{d}\, P_\epsilon(x)$

Holomorphic-Topological Feynman Diagrams

- Non-vanishing Feynman diagrams are **Laman graphs** and generalizations



- Feynman diagrams and integrals satisfy infinite collections of geometric identities relating to the configuration space of almost-Laman graphs
  - Feynman integrals satisfy infinite collection of **quadratic identities**
  - Identities imply (higher)-associativity of the accompanying brackets in a diagram-by-diagram way
  - Can **bootstrap** all Feynman integrals from these identities?
- Can use these Laman graphs to prove a completely combinatorial/diagrammatic proof of **perturbative non-renormalization** of $T \geq 2$ theories.

Recap

I. Reviewed basic examples of deformations in QFT (β-function)
II. Introduced **η-function**; interactions have L_∞-algebra structure, tracks violation of BRST symmetry by interactions
III. η-function contains familiar data like anomaly data, and briefly discussed the relation to twisted SQFTs
IV. Introduced holomorphic-topological theories, and showed brackets are very strongly constrained (**Laman graphs**)
V. Laman graphs prove associativity relations, and no perturbative corrections when $T \geq 2$ topological directions.

Three Punchlines

1. η-vector exists and is computable
2. η-vector contains anomalies, OPEs, and more
3. Non-renormalization theorem for HT theories