Title: Collinear singularities from a double cover of twistor space

Speakers: Seraphim Hsieh Jarov

Collection/Series: Celestial Holography Summer School 2024

Date: July 24, 2024 - 3:35 PM

URL: https://pirsa.org/24070081

Abstract:

Based on an idea of Kevin Costello, I will show how to construct a double cover of the twistor space of $\mathbb{R}^4$, $X = \pi^*(\mathcal{O}(1)\oplus\mathcal{O}(1))\to\Sigma$ where Σ is an (hyper)elliptic curve. I then discuss how holomorphic theories such as BF and Chern-Simons theory on X descend to theories on ordinary twistor space. Once on twistor space, compactifying along the $\mathbb{CP}^1$ direction of twistor space produces a corresponding 4d theory where we can study the algebra of collinear singularities. I will present my calculations which show that this algebra lives on the elliptic curve defining the double cover of twistor space.

Building celestial chiral algebras living on elliptic curves

Seraphim Jarov (sjarov@perimeterinstitute.ca)

Twistor space setup

Recall that $\mathbb{PT} = (\mathcal{O}(1) \oplus \mathcal{O}(1) \to \mathbb{CP}^1) \cong \mathbb{CP}^1 \times \mathbb{R}^4$ and

$$\underbrace{\int_{\mathbb{PT}} \mathrm{tr}(\mathrm{bF}^{0,2}(\mathrm{a}))}_{\text{BF Theory}} \xleftrightarrow{\text{Penrose transform}} \underbrace{\int_{\mathbb{R}^4} \mathrm{tr}(\mathrm{B} \wedge \mathrm{F}(\mathrm{A})_+)}_{\text{SDYM}} .$$

Scattering plane waves in 4d SDYM gives collinear singularities that live on $\mathbb{CP}^1$. Eg.



$$\sim \frac{1}{\langle 12 \rangle} e^{\mathrm{ix}\cdot(\mathrm{p}_1+\mathrm{p}_2)} = \frac{1}{\mathrm{z}_1 - \mathrm{z}_2} e^{\mathrm{ix}\cdot(\mathrm{p}_1+\mathrm{p}_2)}.$$

Elliptic curve setup

Let $w^2 = \Pi_i^n (z - z_i) = H(z)$ define an (hyper)elliptic curve $\Sigma \xrightarrow[2:1]{\pi} \mathbb{CP}^1$.

We can construct

$$X = (\pi^* \mathcal{O}(1) \oplus \pi^* \mathcal{O}(1) \to \Sigma) \cong \Sigma \times \mathbb{R}^4 \xrightarrow[2:1]{} \mathbb{PT}$$

and consider

$$\int_X \beta F^{0,2}(\alpha) \xrightarrow{\pi_*} \int_{\mathbb{PT}} b\bar{\partial} a + \tilde{b}\bar{\partial}\tilde{a} + baa + a\tilde{a}\tilde{b} + b\tilde{a}\tilde{a}H(z).$$

Example: Take $\deg H = 4$ (so $\Sigma = \mathbb{T}^2$). The 4d action is

$$\int_{\mathbb{R}^4} BF(A) + \tilde{\Phi}\Delta\Phi + A^{\alpha}_{\dot{\alpha}} D^{\dot{\alpha}}_{\alpha} \Phi\tilde{\Phi} + B_{\alpha\beta} D_{\dot{\alpha}\delta} \Phi D^{\dot{\alpha}}_{\gamma} \Phi H^{\alpha\beta\delta\gamma} + \ldots$$

a deformation of SDYM with two scalars.

Celestial chiral algebra lives on the elliptic curve

$$\int_{\mathbb{R}^4} \mathrm{BF(A)} + \tilde{\Phi}\Delta\Phi + \mathrm{A}^{\alpha}_{\dot{\alpha}}\mathrm{D}^{\dot{\alpha}}_{\alpha}\Phi\tilde{\Phi} + \mathrm{B}_{\alpha\beta}\mathrm{D}_{\dot{\alpha}\delta}\Phi\mathrm{D}^{\dot{\alpha}}_{\gamma}\Phi\mathrm{H}^{\alpha\beta\delta\gamma} + \ldots$$

Scattering plane waves on the double cover gives diagrams like



$$\sim \frac{\omega(\mathrm{p}_1)\omega(\mathrm{p}_2) + \tilde{\omega}(\mathrm{p}_1)\tilde{\omega}(\mathrm{p}_2)}{\mathrm{z}_1 - \mathrm{z}_2}.$$

Singular only when $\mathrm{z}_1 \to \mathrm{z}_2$ on the same branch of the double cover.

Where $\omega, \tilde{\omega}$ are the even and odd components of the plane wave on X.